arXiv:1506.00046v3 [math.PR] 25 Nov 2016

Ray-Knight representation of flows of branching processes with competition by pruning of Lévy trees

J.Berestycki [*], M.C. Fittipaldi [†] and J.Fontbona[‡]

November 28, 2016

Abstract

We introduce flows of branching processes with competition, which describe the evolution of general continuous state branching populations in which interactions between individuals give rise to a negative density dependence term. This generalizes the logistic branching processes studied by Lambert [17]. Following the approach developed by Dawson and Li [10], we first construct such processes as the solutions of certain flow of stochastic differential equations. We then propose a novel genealogical description for branching processes with competition based on interactive pruning of Lévy-trees, and establish a Ray-Knight representation result for these processes in terms of the local times of suitably pruned forests.

Keywords: continuous state branching processes, competition, stochastic flows, Lévy-tree exploration and local times, Poisson-snake, pruning, Ray-Knight representation.

AMS MSC 2010: 60J25; 60G57; 60J80 .

Contents

1 Introduction and main results **2**

1.1 Continuous state branching processes and their genealogies . . . 2

1.2 Flows of branching processes with competition . . . 4

1.3 Lévy CRT and Ray-Knight Theorem for flows of CSBP . . . 5

1.4 Pruning of Lévy trees . . . 7

1.5 Main results . . . 9

1.6 Plan of the paper . . . 10

1.7 Discussion of related results . . . 10

2 Continuous state branching processes with competition **11**

2.1 Basic properties of the stochastic flow . . . 12

---

[*] Department of Statistics, University of Oxford, 1 South Parks Road, Oxford, OX13TG United Kingdom. E-mail: `julien.berestycki@stats.ox.ac.uk`

[†] Universidad de Buenos Aires, Departamento de Matemáticas, Facultad de Ciencias Exactas y Naturales, Ciudad Universitaria - Pabellón I, Buenos Aires-Argentina. E-mail: `mcfittipaldi@dm.uba.ar`

[‡] Department of Mathematical Engineering and Center for Mathematical Modeling, UMI(2807) UCHILE-CNRS, University of Chile, Casilla 170-3, Correo 3, Santiago-Chile. E-mail: `fontbona@dim-uchile.cl`.

3 Competitive pruning **14**
3.1 A Poisson Lévy-snake 14
3.2 Marking at height-time adapted rates 15
3.3 An operator on adapted intensities 17
3.4 Iterative scheme and fixed point 19

4 A Ray-Knight representation **21**
4.1 Strategy of proof of Theorem 1.3 21
4.2 Grid approximation of the stochastic flow of branching processes with competition 22
4.3 Grid approximation of competitively marked local times 24
4.4 Proof of the Ray-Knight representation 26

5 Consistency of the stochastic flow approximation **27**
5.1 Some auxiliary results 27
5.2 Proof of Proposition 4.1 30

6 Law and consistency of the local time approximation **32**
6.1 Poisson-snake excursions 32
6.2 Proofs of Proposition 4.2 and Lemma 4.2 34
6.3 Proof of Proposition 4.3 37

A Appendix **42**
A.1 Proof of Proposition 1.2 42
A.2 Proof of estimate (5.7) 42
A.3 Proof of Lemma 6.1 43
A.4 Proof of Lemma 6.3 44
A.5 Proof of Lemma 6.5 44
A.6 Lamperti-type representation 47

References **48**

1 Introduction and main results

1.1 Continuous state branching processes and their genealogies

Continuous state branching processes model the evolution of the size of a continuum population in which individuals reproduce and die but do not interact. Mathematically, a continuous state branching process (or CSBP, for short) is a càdlàg $[0,\infty)$-valued strong Markov processes $Y = (Y_t : t \geq 0)$ with law $\mathbb{P}_y$ given the initial state $y \geq 0$, satisfying the branching property: for any $y_1, y_2 \in [0,\infty)$, Y under $\mathbb{P}_{y_1+y_2}$ has the same law as the independent sum $Y^{(1)} + Y^{(2)}$, with $Y^{(i)}$ distributed as Y under $\mathbb{P}_{y_i}$ $(i = 1, 2)$. CSBPs arise as the possible scaling limits of discrete Galton-Watson processes, and their laws are completely characterized by their so-called *branching mechanism*, which is the Laplace exponent $\lambda \mapsto \psi(\lambda)$ of a spectrally positive Lévy process X. More precisely, one has

$$\mathbb{E}_x(e^{-\theta Y_t}) = e^{-x u_t(\theta)},\ x \geq 0,$$

where u_t is the unique nonnegative solution of the differential equation

$$\frac{\partial u_t(\theta)}{\partial t} = -\psi(u_t(\theta)), \quad u_0(\theta) = \theta.$$

By a celebrated result of Lamperti, Y can also be obtained in a pathwise way from X by means a random time-change [18]. We term the process Y a ψ-CSBP to specify the underlying branching mechanism.
In this work, we focus on the case where the function $\psi(\lambda) = \ln \mathbb{E}(e^{-\lambda X_1})$ has the following properties (1.1) and (1.2), which **we assume throughout**:
One has

$$\psi(\lambda) = \alpha\lambda + \frac{1}{2}\sigma^2\lambda^2 + \int_{(0,\infty)} (e^{-\lambda x} - 1 + \lambda x)\Pi(dx)\,, \quad \lambda \geq 0\,, \tag{1.1}$$

for some $\alpha, \sigma \geq 0$ and Π a measure on $(0,\infty)$ such that $\int_{(0,\infty)}(x \wedge x^2)\Pi(dx) < \infty$. Moreover, Grey's condition holds, that is, one has

$$\int_1^\infty \frac{d\lambda}{\psi(\lambda)} < \infty. \tag{1.2}$$

Condition (1.1) ensures that Y is conservative (i.e. $\forall\, t, x > 0$, $\mathbb{P}_x(Y_t < \infty) = 1$) and (sub)critical ($\psi'(0+) \geq 0$). Condition (1.2) implies that there is a.s. extinction (i.e. $\mathbb{P}_x(\exists t \geq 0 : Y_t = 0) = 1$, $\forall$ $x > 0$) and that $\sigma > 0$ or $\int_0^1 r\Pi(dr) = \infty$, so the paths of Y have infinite variation a.s. We refer the reader to [12], [16], [21] for these facts and for general background on CSBP.
The branching property allows us to construct a family of ψ-CSBP as a two parameters process $(Y_t(v), t \geq 0, v \geq 0)$, with v ranging over all possible initial population sizes. In a more general setting, this was first done by Bertoin and Le Gall [8] using families of nested subordinators. More recently, Dawson and Li constructed such a process by means of a *stochastic flow* of SDE driven by Gaussian space-time white noise and a Poisson random measure. Precisely, in [10] the process $(Y_t(v) : t \geq 0, v \geq 0)$ was obtained as the unique strong solution of the family stochastic differential equation:

$$\begin{aligned} Y_t(v) = v - \alpha \int_0^t Y_s(v)\mathrm{d}s + \sigma \int_0^t \int_0^{Y_{s-}(v)} W(\mathrm{d}s, \mathrm{d}u) + \int_0^t \int_0^{Y_{s-}(v)} \int_0^\infty r\tilde{N}(\mathrm{d}s, \mathrm{d}\nu, \mathrm{d}r), \\ t \geq 0, v \geq 0, \end{aligned} \tag{1.3}$$

where $W(\mathrm{d}s, \mathrm{d}u)$ is a Gaussian white noise process on $[0,\infty)^2$ based on the Lebesgue measure $\mathrm{d}s \otimes \mathrm{d}u$ and $\tilde{N}$ is the compensated version of a Poisson random measure N on $[0,\infty)^3$ with intensity $\mathrm{d}s \otimes \mathrm{d}\nu \otimes \Pi(\mathrm{d}r)$. Stochastic calculus easily yields the fact that, for each $v \geq 0$, the above process is a ψ-CSBP, started from the initial population size v. Moreover, it is shown in [10] that $(Y_t(v) : t \geq 0, v \geq 0)$ has a version with the following properties:

i) for each $v \geq 0$, $t \mapsto Y_t(v)$ is a càdlàg process on $[0,\infty)$;

ii) for each $t \geq 0$, $v \mapsto Y_t(v)$ is a non-negative and non-decreasing càdlàg process on $[0,\infty)$;

iii) for any $0 \leq v_1 \leq v_2 \leq \cdots \leq v_n$, the processes $(Y_t(v_j) - Y_t(v_{j-1}) : t \geq 0)$, $j = 1, \ldots n$ are independent CSBP with branching mechanism ψ issued from $v_j - v_{j-1}$ (branching property); and

iv) if for each $0 \leq s$ we denote by $(Y_{s,t}(v) : t \geq s)$ the solution of (1.3) starting at time s, that is,

$$Y_{s,t}(v) = v - \alpha \int_s^t Y_{s,r}(v)\mathrm{d}r + \sigma \int_s^t \int_0^{Y_{s,r-}(v)} W(\mathrm{d}r, \mathrm{d}u) + \int_s^t \int_0^{Y_{s,r-}(v)} \int_0^\infty u\tilde{N}(\mathrm{d}r, \mathrm{d}\nu, \mathrm{d}u), \tag{1.4}$$

then for every $0 \leq s \leq r$ we have a.s. $(Y_{s,t}(v) : t \geq r) = (Y_{r,t}(Y_{s,r}(v)) : t \geq r)$ (flow property).

The family of solutions to equation (1.3) is thus called a *flow of continuous state branching processes* or a *measure-valued branching process* (as in [21]). One upshot of such a construction is that, in a similar way as in [8], one can make sense of the genealogy of the family of CSBPs (1.3): an individual y at time $t \geq 0$ is a descendant of the individual x at time $s \leq t$ if and only if $y \in (Y_{s,t}(x-), Y_{s,t}(x)]$.

There is, however, a more explicit and natural way to encode the genealogy of CSBPs, which relies on tree-like topological random objects known as *continuum random trees* (CRT). This point of view builds upon the pioneer works of Ray [24] and Knight [15] on the quadratic branching case, which imply that the full flow of the Feller diffusion can be constructed from the local times, at different heights, of reflected Brownian motion. Aldous [5, 6] later showed how each of the corresponding Brownian excursions codes a CRT. These results, together with Itô's Poissonian representation of the Brownian excursion process [14] provide in that case a complete description of the genealogical trees of the evolving population. More recently, Le Gall and Le Jan [20] and Duquesne and Le Gall [11] extended such a genealogical representation to a general class of ψ-CSBP. More precisely, under assumptions (1.1) and (1.2), those works provided a *Ray-Knight representation* result for a ψ-CSBP Y in terms of the height and local times processes of certain measure valued Markov process, the *exploration process*, defined using the corresponding Lévy process X reflected at is running infimum (we recall this construction and the precise statement in Section 1.3).

During the last decade, the scope of mathematically tractable population models featuring branching behavior has been considerably enlarged to include models with interactions, immigration and density dependence. Negative density dependence, in particular, can represent competition among individuals, due to limited resources or other mechanisms. The aim of the present work is to extend the Ray-Knight genealogical representation to branching-type populations with competition between individuals.

1.2 Flows of branching processes with competition

The prototypical example of a continuum branching model with competition is the logistic branching process (LPB), introduced by Lambert in [17] by means of a Lamperti transform of spectrally positive Ornstein-Ulhenbeck processes. Alternatively, LBPs are defined as scaling limits of some discrete population models, where the death rate of each individual owed to competition, is proportional to the instantaneous population size. The aggregate competition rate results in that case into a negative drift, proportional to the squared population size (see [17] for details and further discussion).
Generalizing that idea, we consider $g : [0,\infty) \to [0,\infty)$ a locally bounded function, which we refer to as the *competition mechanism*. Heuristically, $g(z)$ is the rate at which an additional individual in a given population of size z would be killed due to competition. We introduce a competitive analogue of (1.3):

Proposition 1.1. *Let* $W(\mathrm{d}s, \mathrm{d}u)$ *and* $N(\mathrm{d}s, \mathrm{d}\nu, \mathrm{d}r)$ *be the same processes considered in* (1.3). *Define* $G(z) := \int_0^z g(u)\mathrm{d}u \geq 0$. *There is, for each* $v \geq 0$, *a unique strong solution of the stochastic differential equation:*

$$\begin{aligned} Z_t(v) = \; & v - \alpha \int_0^t Z_s(v)\mathrm{d}s + \sigma \int_0^t \int_0^{Z_{s-}(v)} W(\mathrm{d}s, \mathrm{d}u) + \int_0^t \int_0^{Z_{s-}(v)} \int_0^\infty r\tilde{N}(\mathrm{d}s, \mathrm{d}\nu, \mathrm{d}r) \\ & - \int_0^t G(Z_s(v))\mathrm{d}s, \quad t \geq 0. \end{aligned} \tag{1.5}$$

Moreover, the process $(Z_t(v) : t \geq 0, v \geq 0)$ *admits a (bi-measurable) version such that*

- *i) for each* $v \geq 0$, $t \mapsto Z_t(v)$ *is a càdlàg process on* $[0,\infty)$*;*
- *ii) for each* $t \geq 0$, $v \mapsto Z_t(v)$ *is a non-negative and non-decreasing càdlàg process on* $[0,\infty)$*;*
- *iii) for each* $u \geq 0$, *the conditional law of* $(Z_t(v) - Z_t(u) : t \geq 0, v \geq u)$ *given* $(Z_t(x) : t \geq 0, 0 \leq x \leq u)$ *depends only on* $(Z_t(u) : t \geq 0)$*; and*

iv) defining for each $s \geq 0$ *the process* $(Z_{s,t}(v) : t \geq s, v \geq 0)$ *solution to*

$$Z_{s,t}(v) = v - \alpha \int_s^t Z_{s,r}(v)\mathrm{d}r + \sigma \int_s^t \int_0^{Z_{s,r-}(v)} W(\mathrm{d}r, \mathrm{d}u) + \int_s^t \int_0^{Z_{s,r-}(v)} \int_0^\infty u\tilde{N}(\mathrm{d}r, \mathrm{d}\nu, \mathrm{d}u) \\ - \int_s^t G(Z_{s,r}(v))\mathrm{d}r,$$

we have for every $0 \leq s \leq r$ *a.s.* $(Z_{s,t}(v) : t \geq r) = (Z_{r,t}(Z_{s,r}(v)) : t \geq r)$.

We call $(Z_t(v) : t \geq 0, v \geq 0)$ a stochastic flow of branching processes with branching mechanism ψ and competition mechanism g, or simply *stochastic flow of branching processes with competition*, if the mechanisms are known and fixed.

In the case that $g = C > 0$ is constant, one clearly recovers a flow of CSBP as in (1.3) but, in general, the branching property is lost (notice the change in property (iii) between solutions of (1.3) and solutions of (1.5)). It is easily checked that in the linear case, when $g(x) = 2cx$ say, for each $v \geq 0$ the process $t \mapsto Z_t(v)$ reduces to the usual logistic branching process studied in [17] (with competition intensity $c > 0$). If moreover $\Pi = 0$, one gets the so-called logistic Feller diffusion

$$\mathrm{d}Z_t = \left(-\alpha Z_t - cZ_t^2\right)\mathrm{d}t + \sigma\sqrt{Z_t}\mathrm{d}B_t, \quad Z_0 = v, \tag{1.6}$$

for which the case $-\alpha \geq 0$ has been studied by Le *et al.* [19] and Pardoux and Wakolbinger [23].

The competition dynamics implicit in equation (1.5) is determined by an ordering of the population. Indeed, if the complete population is identified with the positive half-line, we are implicitly assuming that an individual at $x > 0$ can only be killed "from below", i.e. by the fraction of the population in $[0, x)$, at rate given by $g(x)$. The downward drift $G(Z_t(v)) = \int_0^{Z_t(v)} g(u)\mathrm{d}u$ thus corresponds to the total rate of killing at time t in the population started from size v at time 0. This picture is similar to the point if view adopted in [19, 23] to establish a genealogical representation for the logistic Feller diffusion, which we further discuss later.

Although one can already read the genealogy structure in (1.5), our main aim here is to describe this genealogy using a Lévy CRT, that is, in a manner analogous to the Ray-Knight theorem for CSBP. Following [20] and [11], we next recall in details that result and introduce some basic objects we require.

1.3 Lévy CRT and Ray-Knight Theorem for flows of CSBP

We briefly review here the exploration process, introduced in [20] to construct the Lévy continuum random tree from a spectrally positive Lévy process and used in [11] to define the continuum genealogy of the associated CSBP. We follow [11] and [3] and refer to Chapter 1 of the former for detailed proofs and further background.

Let X be a Lévy process with Laplace exponent ψ satisfying the standing assumptions. Zero is then regular for X reflected at its *running infimum*. We denote the latter process by $I_t := \inf_{0 \leq s \leq t} X_s$ and recall that $-I$ is a local time at 0 for the strong Makov process $X - I$. For $0 \leq s \leq t$ we denote by $I_t^s = \inf_{s \leq r \leq t} X_r$ the two parameter process known as *future infimum* of X. The *height process* $H^0 = (H_t^0 : t \geq 0)$ is defined for each $t \geq 0$ as $H_t^0 = \liminf_{\varepsilon \to 0} \frac{1}{\varepsilon} \int_0^t \mathbf{1}_{\{X_s < I_t^s + \varepsilon\}} ds$ (which, by time reversal, is the total local time of the dual Lévy process on $[0, t]$ reflected at its supremum) and measures for each $t \geq 0$ the size of the set $\{s \leq t : X_s = \inf_{[s,t]} X_r\}$. This process is neither Markovian nor a semimartingale unless X has no jumps (in which case it is a reflected Brownian motion with drift). However it does always have a version which is a measurable function of some strong Markov process, called exploration process. The *exploration process* $\rho = (\rho_t : t \geq 0)$ takes values in the space of finite measures in $\mathbb{R}_+$ and, for each $t \geq 0$, it is defined on bounded measurable functions f by

$$\langle \rho_t, f \rangle = \int_0^t d_s I_t^s f(H_s^0),$$

where $d_s I_t^s$ denotes Lebesgue-Stieltjes integration with respect to the nondecreasing map $s \mapsto I_t^s$. Equivalently, one can write

$$\rho_t(dr) = \beta \mathbf{1}_{[0,H_t^0]}(r)dr \; + \sum_{0<s\leq t, X_{s-}<I_t^s} (I_t^s - X_{s-})\delta_{H_s^0}(dr). \tag{1.7}$$

In particular, the measure ρ_t can be written as a function of the excursion of the reflected Lévy process $(X_s - I_s \,:\, s \geq 0)$ straddling t. Furthermore, $t \mapsto \rho_t$ is càdlàg in the variation norm, ρ_t has total mass $\langle \rho_t, 1 \rangle = X_t - I_t$ for each $t \geq 0$ and $\rho_t(\{0\}) = 0$. The process defined by $H_t := \sup \operatorname{supp}(\rho_t)$ (with $\operatorname{supp}(\mu)$ the topological support of μ and the convention that $\sup \emptyset = 0$) is a continuous modification of H_t^0 and one has $\operatorname{supp}(\rho_t) = [0, H_t]$ when $\rho_t \neq 0$. This also implies that the excursion intervals out of zero are the same for $X - I$, ρ or H. We call $\mathbf{N}$ the excursion measure (away from zero) of the strong Markov process ρ or, equivalently, of $X - I$.

Under $\mathbf{N}$, the process $s \mapsto H_s$ is a.e. continuous, non-negative with compact support and we have $H_0 = 0$. It therefore encodes a tree, as follows. Let $\zeta = \inf\{s > 0 : H_s = 0\}$ denote the length of the canonical excursion under $\mathbf{N}$. The function d_H on $[0,\zeta]^2$ given by

$$d_H(s,t) = H_s + H_t - 2H_{s,t} \quad \text{with} \quad H_{s,t} := \inf_{s\wedge t\leq r\leq s\vee t} H_r\,,$$

defines an equivalence relation $\sim_H$ whereby $(s \sim_H t) \Leftrightarrow (d_H(s,t) = 0)$, $\forall s,t \leq \zeta$. Hence, d_H induces a distance on the quotient space $\mathcal{T}_H = [0,\zeta]/\sim_H$ which is thus a compact metric space, more precisely, a "real tree". That is, any two points are joined by a unique, up to re-parametrization, continuous injective path, isomorphic to a line segment [1].

The *ψ-Lévy random tree* (or Lévy CRT) is the real tree $(\mathcal{T}_H, d_H)$ coded by H under the measure $\mathbf{N}$. Hence, each $s \in [0,\zeta]$ labels a vertex at height H_s in the tree and $d_H(s,t)$ is the distance between vertices corresponding to s and t; accordingly, $H_{s,t}$ represents the height (or generation) of the most recent ancestor common to s and t. The equivalence class of $s = 0$ is termed *root* and the unique path isomorphic to $[0, H_t]$ connecting it with the class of $t \geq 0$, is interpreted as the ancestral line or "lineage" of the individual corresponding to t. Thus, ρ_t can be seen as a measure on this lineage, describing the mass of sub-trees grafted on its right.

Under $\mathbb{P}$, the process of excursions of ρ is Poisson with respect to the local time clock at level 0, with intensity measure $\mathbf{N}$ (much as reflected Brownian motion is a Poisson point process of Brownian excursions). The genealogy of the population is then described by the "Levy forest" $\mathcal{T}$ defined as the union of the corresponding Poissonian collection of trees.

Since $H_t = 0$ if and only if $X_t - I_t = 0$, the local time at level 0 of H is naturally defined as the process $L_t^0 := -I_t$. One way of defining the local time process $(L_t^a : t \geq 0)$ of H at levels $a > 0$ is through the spatial Markov property of the exploration process: if for each $t \geq 0$ we set

$$\tau_t^a := \inf\{s \geq 0 : \int_0^s \mathbf{1}_{\{H_r > a\}} dr > t\} = \inf\{s \geq 0 : \int_0^s \mathbf{1}_{\{\rho_r((a,\infty))>0\}} dr > t\} \tag{1.8}$$

and $\tilde{\tau}_t^a := \inf\{s \geq 0 : \int_0^s \mathbf{1}_{\{H_r \leq a\}} dr > t\}$, then the measure valued process $(\rho_t^a : t \geq 0)$ defined by

$$\langle \rho_t^a, f \rangle = \int_{(a,\infty)} \rho_{\tau_t^a}(dr) f(r-a) \tag{1.9}$$

---

[1] The reader is referred to [12] for further topological background.

has the same distribution as $(\rho_t : t \geq 0)$, and is independent of the sigma field generated by the process $((X_{\tilde{\tau}_t^a}, \rho_{\tilde{\tau}_t^a}) : t \geq 0)$. Thus, for each $t \geq 0$, we define $L_t^a := l^a \left(\int_0^t \mathbf{1}_{\{H_r > a\}} dr \right)$ where $l^a := (l^a(s) : s \geq 0)$ denotes the local time at 0 of the Lévy process reflected at its infimum $(\langle \rho_s^a, 1 \rangle : s \geq 0)$.
The Ray-Knight theorem of Duquesne and Le Gall [11, Theorem 1.4.1] states that, for each $x \geq 0$, a ψ-CSBP starting at x is equal in law to the processes $(L_{T_x}^a : a \geq 0)$, where

$$T_x := \inf\{t \geq 0 : L_t^0 = x\}. \tag{1.10}$$

Thanks to the strong Markov property of the exploration process and to the branching property iii) of process (1.3), this result obviously extends to a two-parameter processes as follows:

Theorem 1.1 (Ray-Knight representation for flows of CSBP). *The process* $(L_{T_x}^a : a \geq 0, x \geq 0)$ *and the process* $(Y_a(x) : a \geq 0, x \geq 0)$ *given by* (1.3) *have the same law.*

1.4 Pruning of Lévy trees

Following [3, 2], the *pruning* of a real tree or forest $\mathcal{T}$ at a discrete of points $\bar{\tau} \subset \mathcal{T}$ is the subset of $\mathcal{T}$ defined as the union of the connected components of $\mathcal{T} \setminus \bar{\tau}$ containing the roots. According to those works, if conditionally on the corresponding Lévy forest $\mathcal{T}$ the point configuration is randomly distributed as a Poisson point process of intensity $\theta > 0$ (with respect to the natural length measure on the skeleton of $\mathcal{T}$), then the resulting pruned random subforest $\mathcal{T}^\theta$ has the same law as the the random Lévy forest associated with the branching mechanism

$$\psi_\theta(\lambda) := \psi(\lambda) + \theta\lambda\,; \tag{1.11}$$

see [3] and Proposition 1.2 below for rigorous statements in terms of exploration and local times processes, respectively. In order to formulate a genealogical representation for process (1.5) analogue to Theorem 1.1, we will extend those ideas, by pruning the Lévy forest at variable rates.
To that end, we need to first formalize the notion of a Poissonian configuration of points on the product space $\mathcal{T} \times \mathbb{R}_+$, with, as intensity, the product of the respective length (Lebesgue) measures. The following notation will be used in the sequel:

- $\mathscr{M}_f^0(\mathbb{R}_+)$ stands for the space of compactly supported Borel measures on $\mathbb{R}_+$. For $\mu \in \mathscr{M}_f^0(\mathbb{R}_+)$, we set $H(\mu) := \sup \operatorname{supp}(\mu)$, where $\operatorname{supp}(\mu)$ is the topological support of μ and $\sup \emptyset := 0$.
- $\mathscr{M}_{at}(\mathbb{R}_+^2)$ denotes the space of atomic Borel measures on $\mathbb{R}_+^2$ with unit mass atoms.
- $\mathcal{V} := \left\{(\mu, \eta) \in \mathscr{M}_f^0(\mathbb{R}_+) \times \mathscr{M}_{at}(\mathbb{R}_+^2) : \ \operatorname{supp} \mu = [0, H(\mu)) \text{ and } \operatorname{supp} \eta \subseteq [0, H(\mu)) \times \mathbb{R}_+\right\}$.

The object next defined is an instance of the snake processes introduced in [11]. It extends, in a way, a Poisson Lévy-snake used in [3] to prune a forest at constant rate, and it is essentially a variant of the objects used in [4, 1] to define fragmentation processes by means of Lévy trees. (See Section 3.1 below for its construction as a snake process and the precise topology that is put in $\mathcal{V}$.)

Definition 1.1. We call exploration process with positive marks or simply *marked exploration process* a càdlàg strong Markov process $((\rho_s, \mathcal{N}_s) : s \geq 0)$ with values in $\mathcal{V}$ such that

1. $(\rho_s : s \geq 0)$ is the exploration process associated with the Lévy process X;
2. conditionally on $(\rho_s : s \geq 0)$, for each $s \geq 0$, $\mathcal{N}_s$ is a Poisson point measure on $[0, H_s) \times \mathbb{R}_+$ with intensity the Lebesgue measure. Moreover, for all $0 \leq s \leq s'$,

- $\mathcal{N}_{s'}(\mathrm{d}r,\mathrm{d}\nu)\mathbf{1}_{\{r\le H_{s,s'}\}}=\mathcal{N}_s(\mathrm{d}r,\mathrm{d}\nu)\mathbf{1}_{\{r\le H_{s,s'}\}}$ a.s. and

- $\mathcal{N}_{s'}(\mathrm{d}r,\mathrm{d}\nu)\mathbf{1}_{\{r> H_{s,s'}\}}$ and $\mathcal{N}_s(\mathrm{d}r,\mathrm{d}\nu)\mathbf{1}_{\{r> H_{s,s'}\}}$ are independent point processes.

The last two properties in point 2. naturally correspond to the random tree structure of $\mathcal{T}$ and are classically referred to as the *snake property*. Thanks to them, the Poisson snake $\mathcal{N}$ induces a point process in the obvious quotient space $\mathcal{T}\times\mathbb{R}_+$ of $\mathbb{R}_+\times\mathbb{R}_+$. Hence, the Poisson process $\mathcal{N}_s$ describes the restriction of that point process to unique path in $\mathcal{T}$ isometric to $[0,H_s]$ joining the root and the point labeled $s\ge 0$.

Note that given $\theta>0$, the point process defined for each $t\ge 0$ by

$$m_t^\theta(A)=\mathcal{N}_t(A\times[0,\theta])\ \forall \text{ Borel set } A\subseteq[0,H_t),$$

is standard Poisson of rate θ on $[0,H_t)$, conditionally on $\mathcal{T}$. The pair $((\rho_t,m_t^\theta):t\ge 0)$ thus corresponds to the exploration process "marked on the skeleton at rate θ" of [3]. According to that work, the "pruned exploration process" defined by $(\rho_t^\theta:t\ge 0):=(\rho_{C_t^\theta}:t\ge 0)$, with C_t^θ the right-continuous inverse of

$$A_t^\theta:=\int_0^t \mathbf{1}_{\{m_s^\theta=0\}}\mathrm{d}s,$$

codes the genealogy of the population of individuals with no marked ancestors (cf. time changing by C_t^θ when exploring the tree amounts to skipping individuals with and atom of m^θ in their lineages). It was then shown in [3, Theorem 0.3] that $(\rho_{C_t^\theta}:t\ge 0)$ has the same law as the exploration process of a CSBP with branching mechanism ψ_θ as in (1.11). That result can be restated in a way that connects pruning, local times and stochastic flows of CSBP, which is relevant to both motivate and prove our main results:

Proposition 1.2. *Let $\theta>0$ be fixed and for each $t,a\ge 0$ define*

$$L_t^a(m^\theta):=\int_0^t \mathbf{1}_{\{m_s^\theta=0\}}\mathrm{d}L_s^a=\int_0^t \mathbf{1}_{\{m_s^\vartheta([0,H_s))=0\}}\mathrm{d}L_s^r.$$

Then, the process defined by $\left(L_{T_x}^a(m^\theta):a\ge 0,x\ge 0\right)$ with T_x as in (1.10)*, has the same law as the stochastic flow of CSBP*

$$\begin{aligned}Y_a^\theta(x)=x-\alpha\int_0^a Y_s^\theta(x)\mathrm{d}s+\sigma\int_0^a\int_0^{Y_{s-}^\theta(x)}W(\mathrm{d}s,\mathrm{d}u)\\ +\int_0^a\int_0^{Y_{s-}^\theta(x)}\int_0^\infty r\tilde{N}(\mathrm{d}s,\mathrm{d}\nu,\mathrm{d}r)-\int_0^a\theta Y_s^\theta(x)\mathrm{d}s,\quad t\ge 0,x\ge 0.\end{aligned}\tag{1.12}$$

The proof of Proposition 1.2 is given in Appendix A.1. In words, this result states that pruning $\mathcal{T}$ at constant rate θ and measuring the local times $\left(L_t^a(m^\theta):t\ge 0,a\ge 0\right)$ in the obtained subtree, amounts to introducing at the level of stochastic flows an additional drift term that corresponds to a constant competition mechanism $g(y)=\theta$. This suggests us that, in order to obtain a pruned forest whose local times equal in law a flow of branching processes with general competition g (as in (1.5)), one should first be able to prune $\mathcal{T}$ at a rate that can dynamically depend on the "previous information" in the exploration-time and height senses.

With this aim, denote by $\mathcal{P}red(\mathcal{N})$ (for predictable) the sigma-field generated by continuous two-parameter processes, whose value at each point (t,h) is measurable with respect to the exploration process ρ up to time t, and to atoms of the Poisson snake $\mathcal{N}$ below height h up to time t (see (3.1) and Definition 3.2 in Section 3.2 for the precise definitions).

Definition 1.2. A real process $(\vartheta(t,h) : h \geq 0, t \geq 0)$ is called a height-time adapted intensity process or simply *adapted intensity process* if it has the following properties:

1. The process $(\vartheta(s,r) : s \geq 0, r \geq 0)$ has a version which is $\mathcal{P}red(\mathcal{N})$-measurable.
2. For each $t \geq 0$, $h \mapsto \vartheta(t,h)$ is a.s. locally integrable.
3. For each pair $s, s' \geq 0$ one a.s. has $\vartheta(s',h) = \vartheta(s,h)$ for $\mathrm{d}h$ a.e. $h \leq H_{s,s'}$.

Property 3. above will be refereed to as the *semi-snake property of adapted intensity processes*. Using these objects and the Poisson snake, we can put marks on the tree at height-time adapted rates as follows:

Definition 1.3. Let $((\rho_s, \mathcal{N}_s) : s \geq 0)$ be the marked exploration process and $\vartheta = (\vartheta(t,h) : t \geq 0, h \geq 0)$ be an adapted intensity process. The *exploration process marked at adapted intensity* ϑ is the process $((\rho_s, m_s^{\vartheta}) : s \geq 0)$, with $m_t^{\vartheta} \in \mathscr{M}_{at}(\mathbb{R}_+)$ the point process given for each $t \geq 0$ by

$$m_t^{\vartheta}(A) = \mathcal{N}_t(\{(h,\nu) : h \in A, \nu \leq \vartheta(h,t)\}) \text{ for all Borels set } A \subseteq [0, H_t). \tag{1.13}$$

Last, in order to measure at each height the mass of individuals in the pruned forest up to a given exploration time (generalizing the idea in Proposition 1.2), we introduce:

Definition 1.4. We call local time process pruned at rate ϑ, or m^{ϑ}*-pruned local time*, the process $(L_t^r(m^{\vartheta}) : t \geq 0, r \geq 0)$ defined by

$$L_t^r(m^{\vartheta}) := \int_0^t \mathbf{1}_{\{m_s^{\vartheta}=0\}} \mathrm{d}L_s^r = \int_0^t \mathbf{1}_{\{m_s^{\vartheta}([0,H_s))=0\}} \mathrm{d}L_s^r\,, \quad r \geq 0, t \geq 0.$$

With these notions in hand, we now turn to the questions of what adapted intensity process we should look for, and state our main results.

1.5 Main results

A closer comparison of (1.12) and (1.5) indicates that, in order to obtain a subforest $\mathcal{T}^*$ encoding the genealogy of the process (1.5), we should prune $\mathcal{T}$ at each point at rate equal to g of the local time left of it in $\mathcal{T}^*$. In terms of the objects we have just introduced, this means that the corresponding adapted intensity ϑ^* should solve, in a certain sense, some fixed point equation. The next result gives a precise meaning to such an object and grants its existence:

Theorem 1.2. *Let g be a competition mechanism and let $F : \vartheta \mapsto F(\vartheta)$ denote the operator acting on adapted intensity processes by*

$$F(\vartheta) = \vartheta'\,, \text{ with } \vartheta'(t,h) = g(L_t^h(m^{\vartheta})) \quad \text{for all } (t,h) \in \mathbb{R}_+^2\,.$$

Then, $F(\vartheta)$ is also and adapted intensity process. Moreover, if we assume that

$$\begin{gathered} g : \mathbb{R}_+ \to \mathbb{R}_+ \text{ is non decreasing and for each } M > 0 \text{ there is } c(M) > 0 \text{ such that} \\ |g(x) - g(y)| \leq c(M)|x - y| \quad \forall\, 0 \leq x, y \leq M, \end{gathered} \tag{\textbf{H}}$$

the operator F has a fixed point $\vartheta^*(t,h)$, a.e unique with respect to $\mathbb{P}(d\omega) \otimes \mathrm{d}t \otimes \mathbf{1}_{[0,H_t(\omega))}(h)\mathrm{d}h$.

The following theorem provides the genealogical representation of a flow of branching process with competition we are looking for and is the main result of the paper (note that T_x is the same as in (1.10)).

Theorem 1.3. *Suppose that condition* (**H**) *on the competition mechanism g hold. Let ϑ^* denote the adapted intensity process given by Theorem 1.2. Then, the processes $(L_{T_x}^a(m^{\vartheta^*}) : a \geq 0, x \geq 0)$ and $(Z_a(x) : a \geq 0, x \geq 0)$ given by* (1.5) *are equal in law.*

1.6 Plan of the paper

In Section **2**, we prove Proposition 1.1, using variants of techniques introduced in [10] and some results established therein. A Lamperti type representation for process (1.1) (for fixed v) is also discussed.
In Section **3** we will first settle some technical issues required to prove Theorems 1.2 and 1.3, concerning in particular filtrations and measurability aspects, as well as properties of the operator F. We will then study the sequence of adapted intensities defined by $\vartheta^{(0)} \equiv 0$ and $\vartheta^{(n+1)} = F(\vartheta^{(n)})$, together with the pruned local times they induce. Heuristically, noting that $\vartheta^{(1)}$ yields a pruned local time process which is smaller than the (non pruned) one associated with $\vartheta^{(0)}$, the fact that g is nondecreasing implies that $\vartheta^{(2)}$ is smaller than $\vartheta^{(1)}$. Pruning according to $\vartheta^{(2)}$ then implies "pruning less" than when using $\vartheta^{(1)}$ and this yields that $\vartheta^{(2)} \leq \vartheta^{(3)} \leq \vartheta^{(1)}$. By iterating this reasoning one gets that $\vartheta^{(2)} \leq \vartheta^{(4)} \leq \ldots \leq \vartheta^{(5)} \leq \vartheta^{(3)} \leq \vartheta^{(1)}$. The proof of Theorem 1.2 makes this remark rigorous and is achieved showing that the sequences $(\vartheta^{(2n)})$ and $(\vartheta^{(2n+1)})$ converge to the same limit, which is a fixed point of F.
In Section **4** we will prove Theorem 1.3 by putting in place a careful approximation argument inspired by Proposition 1.2. More precisely, we will first construct an approximation of the stochastic flow of branching processes with competition (1.5) by means of a flow of CSBP with "piecewise frozen" killing rate, specified by constant negative drifts in a rectangular bi-dimensional grid of constant time-step length and initial population size. We will then construct an approximation of the pruned local time process given by Theorem 1.2, pruning the original forest at constant rates inside blocks of some height-local time grid. The blocks of this second grid are defined in such a way that they are in correspondence, *via* Proposition 1.2, with the rectangles of the stochastic flow grid. The proof of Theorem 1.3 will then be deduced, noting first that these two grid approximations globally have (as two parameter processes defined in different spaces) the same law, and then showing that when the rectangle sizes go to 0, each of them does actually converge (in a strong enough way) to the respective processes they intend to approximate. The technical proofs of these facts are differed to the last two sections.
In Section **5** we prove the convergence of the stochastic flow grid-approximation, using stochastic calculus and some technical ideas adapted from [10]. In Section **6**, combining Proposition 1.2 with global properties of stochastic flows and of local times, we first identify the law of the local time grid approximation with that of the stochastic flow grid approximation. Then, we prove the convergence of the former to the pruned local time process constructed in Theorem 1.2, using mainly techniques from snake-excursion theory. Finally, some technical or auxiliary results will be proved in the Appendix.

Before delving in the proofs, we devote the next subsection to a discussion of our model and results and of their relations to related works in the literature.

1.7 Discussion of related results

In [19], (see also [22, 23]) a Ray-Knight representation theorem for the logistic Feller diffusion was obtained. These authors constructed a process H which can roughly be described as a reflected Brownian motion with a negative drift, proportional to the amount of local time to the left and at the same height of the current state. More precisely, H is defined as the (unique in law) solution of the SDE

$$H_s = \frac{2}{\sigma} B_s + \frac{1}{2} L_s^0(H) - \frac{2\alpha}{\sigma} s - c \int_0^s L_r^{H_r}(H) \mathrm{d}r, \quad s \geq 0, \tag{1.14}$$

where $\alpha < 0$, B is a standard Brownian motion and $L_s^a(H)$ the local time accumulated by H at level $a \geq 0$ up to time $s \geq 0$.
The main result is that the local time process of H, i.e. the process $((\sigma^2/4) L_{T_x}^a(H) : a \geq 0)$, is a weak solution of (1.6) (here, T_x is the stopping time $T_x = \inf\{s > 0 : L_s^0(H) > x\}$ which is unchanged if H is replaced by B). The downward drift of H can thus be understood as a rate of killing which increases

the farther to the right one looks, so that excursions appearing later with respect to exploration time tend to be smaller. Individuals are thus represented as being arranged “from left to right” and as though interacting through “pairwise fights” which are always won by the individual “to the left” (hence lethal for the individual “to the right”). Under another (more pathwise) guise, we adopt the same “trees under attack” picture here using pruning (notice also that solutions of (1.5) generalize a path-valued Markov process used in [19]). There is nevertheless no obvious extension, to the present setting, of the arguments and techniques in [19], which strongly relied on the SDE description (1.14) of the corresponding height process. We must point out however that our results do not cover the case $\alpha < 0$ (hence, a supercritical branching mechanism) treated therein.

In [7], Berestycki et al. study measure valued branching processes *with interactive immigration*, as introduced and discussed by Li in [21], in the spirit of the systems of stochastic differential equations 1.3 of [10]. The main result in [7] is a Ray-Knight representation for those branching processes with immigration. More precisely, suppose that we want a representation of the solutions of

$$\begin{cases} X_t(v) = v + \int_0^t \int_0^{X_{s-}(v)} \int_0^\infty r \tilde{N}(\mathrm{d}s, \mathrm{d}\nu, \mathrm{d}r) + v \int_0^t g(X_s(1))\, ds, \\ v \in [0,1],\ t \geq 0, \end{cases} \tag{1.15}$$

where g is now an *immigration* mechanism, supposed to be non-negative, monotone non-decreasing, continuous and locally Lipschitz continuous away from zero. Observe that this is very similar to equation (1.5) with $\sigma = 0$ and a positive drift term instead of a negative one, which is however also dependent on the current state of the population. Let $(s_i, u_i, a_i, \mathbf{e}_i)$ be a Poisson point process with intensity measure $\mathrm{d}s \otimes \mathrm{d}u \otimes \mathrm{d}a \otimes \mathbf{N}(\mathrm{d}(\mathbf{e}))$ (with $\mathbf{N}$ denoting here the excursion measure of the height process H) and define w_t^i to be the total local time at level t of the excursion $\mathbf{e}_i$. It is shown in [7] that, for all $v \in [0,1]$, there is a unique càdlàg process $(Z_t(v), t \geq 0)$ satisfying $\mathbb{P}$-a.s.

$$\begin{cases} Z_t(v) = \sum_{s_i > 0} w_t^i \, \mathbb{1}_{(a_i \leq v)} \mathbb{1}_{(u_i \leq g(Z_{s_i -}(1)))}, & t > 0, \\ Z_0(v) = v. \end{cases} \tag{1.16}$$

and Z is a weak solution of (1.15).[2] The existence and uniqueness of the process Z is analogue to our Theorem 1.2 (and is also proved by a fixed point argument) while the fact that Z is a solution of (1.15) is analogue to our Theorem 1.3. Let us underline, however, that in [7] the positive drift represents an interactive *immigration* while here we have a negative drift corresponding to a *competition*. This is reflected of course in the fact that instead of *adding* excursions as in [7] we must here *prune* the real forest coding the evolution of the population. In spite of the similarities, adapting techniques in [7] to the present setting does not seem to be simple.

2 Continuous state branching processes with competition

Recall that $G(x) = \int_0^x g(y)\mathrm{d}y$ denotes the primitive of the competition mechanism g. In this section, given a fixed function $z \in \mathbb{D}(\mathbb{R}_+, \mathbb{R}_+)$, we will moreover consider a time inhomogeneous competition mechanism $g_z : \mathbb{R}_+ \times \mathbb{R}_+ \to \mathbb{R}_+$ defined as $g_z(t, y) = g(z_t + y)$. We denote its primitive in v by $G_z(t, x) := \int_0^x g(z_t + y)\mathrm{d}y$, which is locally Lipschitz in x, locally uniformly in t.

Also, we will repeatedly use the following simple result (analogue to Lévy’s characterization of n-dimensional Brownian motion) the proof of which is omitted:

---

[2] The full result is actually that the Poisson point process can be constructed to obtain a strong solution with respect to a given noise to equation (1.15).

Lemma 2.1. *Given* $(\mathcal{S}_t)_{t\geq 0}$ *a filtration in some probability space and* $n, m \in \mathbb{N}$, *let* $(M_t^i : t \geq 0)_{i\leq n}$ *and* $(N_t^k : t \geq 0)_{k\leq m}$ *respectively be continuous* $(\mathcal{S}_t)$*-local martingales and* $(\mathcal{S}_t)$*-adapted counting processes such that, for some real numbers* $a_i > 0, i \leq n$, *and* $b_k > 0, k \leq m$, *one has:*

i) for every $i, j \leq n$, $\langle M^i, M^j\rangle_t = a_i \delta_{ij} t$ *and all* $t \geq 0$ *(with* δ_{ij} *the Kronecker delta); and*

ii) for every $k, l \leq m$ *the processes* $(N_t^k)_{t\geq 0}$ *and* $(N_t^l)_{t\geq 0}$ *have no simultaneous jumps, and* $(N_t^k)_{t\geq 0}$ *has the predictable compensator* $b_k t$.

Then, $(a_i^{-1/2} M_t^i : t \geq 0)_{i\leq n}$ *are standard* $(\mathcal{S}_t)$*-Brownian motions,* $(N_t^k : t \geq 0)_{k\leq m}$ *are standard* $(\mathcal{S}_t)$*-Poisson processes with respective parameters* $(b_k)_{k\leq m}$ *and all these processes are independent.*

2.1 Basic properties of the stochastic flow

Proof of Proposition 1.1 . We first prove parts i) and iii), leaving ii) and iv) for the end of the proof.
i) The tuple of functions (b, σ, g_0, g_1) defined by

- $x \mapsto b(x) := -\alpha x - G(x)$;
- $(x, u) \mapsto \sigma(x, u) := \sigma \mathbf{1}_{\{u\leq x\}}$;
- $(x, \nu, r) \mapsto g_0(x, \nu, r) = g_1(x, \nu, r) := r\mathbf{1}_{\{\nu\leq x\}}$,

are admissible parameters in the sense of conditions $(2.a, b, c, e)$ in [10, Section 2]. Then, [10, Theorem. 2.5] readily provides for each $v \geq 0$ the existence of a unique strong solution to (1.5) and proves i).
iii) Observe first that for each $v \geq 0$ and each $z \in \mathbb{D}(\mathbb{R}_+, \mathbb{R}_+)$ the same arguments of [10, Theorem. 2.5] can be used to prove strong existence and pathwise uniqueness for the stochastic differential equation

$$\begin{aligned} Z_t^z(v) &= v - z_0 - \alpha \int_0^t Z_s^z(v)\mathrm{d}s + \sigma \int_0^t \int_0^{Z_{s-}^z(v)} W(\mathrm{d}s, \mathrm{d}u) \\ &\quad + \int_0^t \int_0^{Z_{s-}^z(v)} \int_0^\infty r\tilde{N}(\mathrm{d}s, \mathrm{d}\nu, \mathrm{d}r) - \int_0^t G_z(s, Z_s^z(v))\mathrm{d}s, \quad t \geq 0. \end{aligned} \tag{2.1}$$

In particular, the solution $Z_t^z(v)$ is adapted to the filtration generated by W and N. We claim that, for each fixed $v \geq u \geq 0$ the process $\Upsilon_t := Z_t(v) - Z_t(u)$, $t \geq 0$ satisfies a similar equation, but with a randomized z. Indeed, from (1.5) we have

$$\begin{aligned} \Upsilon_t &= v - u - \alpha \int_0^t \Upsilon_s \mathrm{d}s + \sigma \int_0^t \int_0^{\Upsilon_{s-}} W'(\mathrm{d}s, \mathrm{d}w) \\ &\quad + \int_0^t \int_0^{\Upsilon_{s-}} \int_0^\infty r\tilde{N}'(\mathrm{d}s, \mathrm{d}\nu, \mathrm{d}r) - \int_0^t G_{Z.(u)}(s, \Upsilon_s)\mathrm{d}s, \quad t \geq 0\,, \end{aligned} \tag{2.2}$$

where now

$$W'(\mathrm{d}s, \mathrm{d}w) = W(\mathrm{d}s, \mathrm{d}w + Z_{s-}(u))$$

is an orthogonal martingale measure (in the sense of Walsh [25]) and

$$N'(\mathrm{d}s, \mathrm{d}\nu, \mathrm{d}r) = N(\mathrm{d}s, \mathrm{d}\nu + Z_{s-}(u), \mathrm{d}r)$$

is a random point measure. Using standard properties of integration with respect to W and N, we can then check that for arbitrary Borel sets $A_1, \ldots A_n$ in $\mathbb{R}_+$ and $B_1, \ldots, B_m$ in $\mathbb{R}_+^2$, disjoint and of finite Lebesgue measure, the processes $M_t^i := W'([0, t] \times A_i)$ and $N_t^k := N'([0, t] \times B_k)$ satisfy the assumptions of Lemma 2.1 with the filtration given by $\mathcal{S}_t := \sigma\left(W([0, s], \mathrm{d}w), N([0, s], \mathrm{d}r, \mathrm{d}\nu), s \leq t\right)$

and with constants $a_i := \int_{A_i} \mathrm{d}w < \infty$ and $b_k = \int \int_{B_k} \Pi(r)\mathrm{d}r\mathrm{d}\nu < \infty$. This implies that W' and N' are respectively a Gaussian white noise process with intensity $\mathrm{d}s \otimes \mathrm{d}w$ and a Poisson random measure with intensity $\mathrm{d}s \otimes \mathrm{d}\nu \otimes \Pi(\mathrm{d}r)$, both with respect to $(\mathcal{S}_t)_{t\geq 0}$, and independent of each other.

In order to prove iii), it is thus enough to show that the pair (W', N') is independent from $\mathcal{Z}^u := \sigma((Z_t(x) : t \geq 0, 0 \leq x \leq u))$ (indeed, we can then apply pathwise uniqueness for (2.1) conditionally on $\mathcal{Z}^u$ to get that Υ is a measurable function of $(W', N', Z(u))$, and immediately deduce the desired property). To that end, we enlarge the probability space with an independent pair $(\hat{W}, \hat{N})$ equal in law to (W, N) and define, given $0 \leq u_1 \leq \cdots \leq u_p \leq u$, the processes

$$\overline{W}(\mathrm{d}s, \mathrm{d}\nu) := \mathbf{1}_{[0,\tilde{Z}_{s-}]}(\nu) W(\mathrm{d}s, \mathrm{d}\nu) + \mathbf{1}_{(\tilde{Z}_{s-},\infty)}(\nu) \hat{W}(\mathrm{d}s, \mathrm{d}\nu - \tilde{Z}_{s-}) \text{ and}$$

$$\overline{N}(\mathrm{d}s, \mathrm{d}\nu, \mathrm{d}r) := \mathbf{1}_{[0,\tilde{Z}_{s-}]}(\nu) N(\mathrm{d}s, \mathrm{d}\nu, \mathrm{d}r) + \mathbf{1}_{(\tilde{Z}_{s-},\infty)}(\nu) \hat{N}(\mathrm{d}s, \mathrm{d}\nu - \tilde{Z}_{s-}, \mathrm{d}r)$$

where $\tilde{Z}_s = \max\{Z_s(u_l) : l = 1, ..., p\}$. Call $(\mathcal{S}^*_t)$ the filtration generated by $(W, \hat{W}, N, \hat{N})$ and, given families of sets $(\overline{A}_i)_{i\leq n'}$ and $(\overline{B}_k)_{k\leq m'}$ with similar properties as $(A_i)_{i\leq n}$ and $(B_k)_{k\leq m}$ considered above, define $\overline{M}^i_t := \overline{W}([0,t] \times \overline{A}_i)$ and $\overline{N}^i_t := \overline{N}([0,t] \times \overline{B}_i)$. Thanks to Lemma 2.1 now applied to the families $\left((M^i)_{i\leq n}, (\overline{M}^i)_{i\leq n'}\right)$ and $\left((N^k)_{k\leq m}, (\overline{N}^k)_{k\leq m'}\right)$ and the filtration $(\mathcal{S}^*_t)$, we readily get that the process $(\overline{W}, \overline{N})$ has the same law as (W, N) and moreover, that (W', N') and $(\overline{W}, \overline{N})$ are independent. Since the processes $Z(u_l), l = 1, ..., p$ are adapted to $(\mathcal{S}^*_t)$ and they also solve a system of SDEs as in (1.5) but driven by $(\overline{W}, \overline{N})$, strong existence and pathwise uniqueness for such equations imply they are measurable functions of $(\overline{W}, \overline{N})$. Hence, their independence from (W', N') follows.

ii) For each $t \geq 0$, the fact that $v \mapsto Z_t(v)$ is increasing can be deduced from the comparison property stated in [10, Thm. 2.3]. Moreover, it is easy to show using similar arguments as in the proof of [10, Prop. 3.4], that there is a locally bounded non-negative function $t \mapsto C(t)$ on $[0, \infty)$ such that

$$\mathbb{E}\left\{\sup_{0\leq s\leq t} |Z_s(v) - Z_s(u)|\right\} \leq C(t)\left\{(v - u) + \sqrt{v - u}\right\} \tag{2.3}$$

for $v \geq u \geq 0$. Alternatively, one can use a simple extension of [10, Thm. 2.3] to stochastic flows with time inhomogeneous drift terms in order to show that, for each $v \geq 0$, the solution $(Z^z_t(v) : t \geq 0)$ of (2.1) when $z_0 = u$ is a.s. dominated on $[0, \infty)$ by the solution $(Y_t(v - u) : t \geq 0)$ of (1.3), then deduce by conditioning first on $(Z_t(x) : t \geq 0, 0 \leq x \leq u)$ (and using part iii)) that

$$\mathbb{E}\left\{\sup_{0\leq s\leq t} |Z_s(v) - Z_s(u)|\right\} \leq \mathbb{E}\left\{\sup_{0\leq s\leq t} |Y'_s(v - u)|\right\},$$

where Y' is defined in terms of (W', N') in the same way as Y was in terms of (W, N), and conclude the bound (2.3) from [10, Prop. 3.4]. Thanks to (2.3) and [10, Lemma 3.5], we can follow the arguments in the proof of [10, Theorem 3.6] to deduce that the path-valued process $(Z(v) : v \geq 0)$ has a càdlàg modification, using when necessary the Markov property stated in iii) instead of the independent increments property of the process $(Y(v) : v \geq 0)$.

iv) This part follows by similar arguments as in Theorem 3.7 of [10]. □

Remark 2.1. Recall that in [17], logistic branching processes were constructed by a Lamperti time-change of certain Lévy-driven Ornstein-Ulhenbeck processes. A similar approach also works here: for a fixed initial population $v > 0$ it is possible to construct the process $Z(v)$ by a Lamperti transform of the solution U to the Lévy-driven SDE

$$\mathrm{d}U_t = \mathrm{d}X_t - \frac{G(|U_t|)}{U_t}\mathrm{d}t, \quad t \geq 0, \quad U_0 = v. \tag{2.4}$$

A precise statement (not required in the sequel) and a sketch of its proof are given in Appendix A.6.

3 Competitive pruning

3.1 A Poisson Lévy-snake

In order to settle some basic properties of the exploration process marked at height-time adapted rate $((\rho_s, m_s^{\vartheta}) : s \geq 0)$ (cf. Def. 1.3), we need to first make precise the way the marked exploration process $((\rho_s, \mathcal{N}_s) : s \geq 0)$ of Definition 1.1 enters into the general framework of Lévy-snake processes constructed in [11, Chapter 4].

Recall that, given (the law of) a Markov process $(\xi_r)_{r\geq 0}$ with càdlàg paths and values in a Polish space E, which is furthermore continuous in probability, the snake with path-space component ξ is a Markov process $((\rho_t, \xi^{(t)}) : t \geq 0)$ with ρ the exploration process and such that, conditionally on ρ, for each $t \geq 0$, $\xi^{(t)} = (\xi_r^{(t)}, r \leq H_t)$ is a path of ξ killed at H_t and for $t < s$, the Markovian paths $\xi^{(t)}$ and $\xi^{(s)}$ are related by the snake property. That is, $\xi^{(t)}$ and $\xi^{(s)}$ are a.s. equal on $[0, H_{t,s}]$ and, conditionally on ρ and one their value at time $H_{t,s}$, their evolutions thereafter are independent (see [11, Proposition 4.1.1]).

For $j = 1, 2$, let us respectively denote by $\mathscr{M}(\mathbb{R}_+^j)$ and $\mathscr{M}_{at}(\mathbb{R}_+^j)$ the space of Borel measures in $\mathbb{R}_+^j$ and the corresponding subspace of atomic Borel measures with unit mass atoms. We endow the space $\mathscr{M}(\mathbb{R}_+)$ with a complete metric inducing the topology of vague convergence (equivalent to weak convergence of the restrictions to every compact subset of $\mathbb{R}_+$). Given π a Poisson point measure on $\mathbb{R}_+^2$ with intensity $\mathrm{d}x \otimes \mathrm{d}y$, seen as a random element of $\mathscr{M}_{at}(\mathbb{R}_+^2)$, we consider the increasing $\mathscr{M}_{at}(\mathbb{R}_+)$-valued additive Markov process $(\xi_r)_{r\geq 0}$ (issued from 0) defined by

$$\xi_r(\cdot) = \pi([0, r] \times \cdot),\ r \geq 0.$$

Then, ξ clearly has the above dicussed properties and we can therefore construct the associated Lévy snake $((\rho_t, \xi^{(t)}) : t \geq 0)$. The marked exploration process $((\rho_t, \mathcal{N}_t) : t \geq 0)$ is then defined setting for each $t \geq 0$:

$$\mathcal{N}_t((a, b] \times A) = \xi_b^{(t)}(A) - \xi_a^{(t)}(A), \forall\, 0 \leq a < b < H_t \text{ and } A \subseteq \mathbb{R}_+ \text{ Borel}$$

and $\mathcal{N}_t(\{0\} \times \cdot) = 0$. This uniquely determines a random element or point process $\mathcal{N}_t \in \mathscr{M}_{at}(\mathbb{R}_+^2)$. It is easy to see that $((\rho_t, \mathcal{N}_t) : t \geq 0)$ thus defined fulfills the distributional conditions in Definition 1.1.

The space $\mathcal{V}$ introduced in Section 1.4 is given a topology as follows. The set $\mathscr{M}_f^0(\mathbb{R}_+)$ is seen as a subset of the space $\mathscr{M}_f(\mathbb{R}_+)$ of finite Borel measures, endowed with the weak topology and an associated complete distance D. Consistently with the previous construction, given a pair $(\mu, \eta) \in \mathcal{V}$ we identify η with the increasing and killed càdlàg path $[0, H(\mu)) \to \mathscr{M}(\mathbb{R}_+)$ given by

$$r \mapsto \eta([0, r] \times \cdot).$$

Finally, as in [11, Section 4.1], we endow the space $\mathcal{V}$ with the complete distance

$$d\left((\mu, \eta), (\mu', \eta')\right) = D(\mu, \mu') + \int_0^{H(\mu)\wedge H(\mu')} \left(d_u(\eta_{(u)}, \eta'_{(u)}) \wedge 1\right) \mathrm{d}u + |H(\mu) - H(\mu')| + d_0(\eta_{(0)}, \eta'_{(0)}),$$

for all $(\mu, \eta), (\mu', \eta') \in \mathcal{V}$, where for each $u \in [0, H(\mu))$, d_u is the Skorohod metric on $\mathbb{D}([0, u], \mathscr{M}(\mathbb{R}_+))$ and $\eta_{(u)}$ is the restriction to $[0, u]$ of the increasing path associated with η [3]. The fact that $((\rho_s, \mathcal{N}_s) : s \geq 0)$ is a càdlàg strong Markov process in $\mathcal{V}$ is then granted by the general results in [11, Chapter 4].

---

[3] Notice that d_0 corresponds to the metric on $\mathscr{M}(\mathbb{R}_+)$.

3.2 Marking at height-time adapted rates

In all the sequel, we will write $(\mathcal{F}_t)_{t\geq 0}$ for the right continuous complete filtration generated by $(\rho_t, \mathcal{N}_t) : t \geq 0)$, and $(\mathcal{F}^{\rho}_t)_{t\geq 0}$ for the one generated by $(\rho_t : t \geq 0)$. For each $t \geq 0$, we moreover introduce $(\mathcal{G}^{(t)}_r)_{r\geq 0}$ the right continuous completion of the filtration given by

$$\sigma\left(\mathcal{F}^{\rho}_t, \{\mathcal{N}_s|_{[0,r]\times\mathbb{R}_+}, s \leq t\}\right), \quad r \geq 0. \tag{3.1}$$

In particular, for each $r, t \geq 0$ we have

$$\mathcal{F}^{\rho}_t \subseteq \mathcal{G}^{(t)}_r \subseteq \mathcal{F}_t.$$

Remark 3.1. Heuristically, $\mathcal{G}^{(t)}_r$ contains all the information to the left of t for ρ and to the left of t and below r for $\mathcal{N}$. Notice that conditionally on $\mathcal{G}^{(t)}_0 = \mathcal{F}^{\rho}_t$, $\mathcal{N}_t$ is a Poisson process in the filtration $(\sigma\left(\mathcal{F}^{\rho}_t, \{\mathcal{N}_t|_{[0,r]\times\mathbb{R}_+}\}\right))_{r\geq 0}$. Hence, $(\mathcal{G}^{(t)}_r)_{r\geq 0}$ is obtained by progressively enlarging that filtration with the sigma-fields

$$\sigma\left(\mathcal{F}^{\rho}_t, \{\mathcal{N}_s|_{[H_{s,t},r]\times\mathbb{R}_+}\mathbf{1}_{\{H_{s,t}<r\}} : s \leq t\}\right), \quad r \geq 0$$

which are independent from it conditionally on $\mathcal{G}^{(t)}_0$, by the snake property of $((\rho_s, \mathcal{N}_s) : s \geq 0)$ (cf. they contain information of marks which are not in the lineage of t).

The following property is then easily deduced:

Lemma 3.1. *For each $t \geq 0$, conditionally on $\mathcal{G}^{(t)}_0$ the point process $\mathcal{N}_t$ on $\mathbb{R}^2_+$ is Poisson of intensity $\mathbf{1}_{[0,H_t)}(r)\mathrm{d}r \otimes \mathrm{d}\nu$, with respect to the filtration $(\mathcal{G}^{(t)}_r)_{r\geq 0}$.*

The use of Poisson calculus for $\mathcal{N}_t$ with respect to the filtration $(\mathcal{G}^{(t)}_r)_{r\geq 0}$ is thus possible but some care is needed regarding measurability issues. We need to introduce

Definition 3.1. For each $t \geq 0$, $\mathcal{P}red^{(t)}$ denotes the sub sigma-field of $\mathcal{B}(\mathbb{R}_+) \otimes \mathcal{F}_t$ generated by $(\mathcal{G}^{(t)}_r)_{r\geq 0}$-adapted processes $(a(r) : r \geq 0)$ which have continuous trajectories.

Definition 3.2. We denote by $\mathcal{P}red(\mathcal{N})$ the sub sigma-field of $\mathcal{B}(\mathbb{R}_+) \otimes \mathcal{B}(\mathbb{R}_+) \otimes \mathcal{F}_\infty$ generated by processes $(b(s,r) : s \geq 0, r \geq 0)$ such that

1. for each $(s,r) \in \mathbb{R}^2_+$, $b(s,r)$ is $\mathcal{G}^{(s)}_r$ measurable,
2. the process $(s,r) \mapsto b(s,r)$ is continuous.

Remark 3.2. A monotone class argument shows that if $(b(s,r) : s \geq 0, r \geq 0)$ is $\mathcal{P}red(\mathcal{N})$- measurable, then $(b(t,r) : r \geq 0)$ is $\mathcal{P}red^{(t)}$-measurable for each $t \geq 0$ (the converse is not true).

Recall that, given an adapted intensity process ϑ, in Definition 1.3 we introduced for each $s \geq 0$ the point process m^{ϑ}_s in $\mathscr{M}_{at}(\mathbb{R}_+)$ defined by

$$m^{\vartheta}_s(A) = \mathcal{N}_s(\{(h,u) : u \leq \vartheta(h,s) \text{ and } h \in A\}), \forall \text{ real Borel set } A.$$

Thus, $((\rho_s, m^{\vartheta}_s) : s \geq 0)$ takes its values in the space of "marked finite measures"

$$\mathbb{S} := \left\{(\mu, w) \in \mathscr{M}^0_f(\mathbb{R}_+) \times \mathscr{M}_{at}(\mathbb{R}_+) : \ \operatorname{supp}(w) \subset [0, H(\mu))\right\}$$

considered in [3]. In a similar way as for elements of $\mathcal{V}$, for $(\mu, w) \in \mathbb{S}$ we now identify the measure w with the increasing element of $\mathbb{D}(\mathbb{R}_+, \mathbb{R})$ given by the path $v \mapsto w([0, v])$ killed at $H(\mu)$ (i.e. the cumulative distribution of w). Following again [11, Chapter 4], $\mathbb{S}$ is endowed with the complete metric

$$\hat{d}((\mu, w), (\mu', w')) := D(\mu, \mu') + \int_0^{H(\mu)\wedge H(\mu')} \left(\hat{d}_u(w_{(u)}, w'_{(u)}) \wedge 1\right) \mathrm{d}u + |H(\mu) - H(\mu')| + \hat{d}_0(w_{(0)}, w'_{(0)}), \tag{3.2}$$

with $\hat{d}_u$ the Skorohod metric on $\mathbb{D}([0, u], \mathbb{R}_+)$ and $w_{(u)}$ the restriction to $[0, u]$ of w.
Next result gathers basic properties of the process $((\rho_t, m_t^\vartheta) : t \geq 0)$ needed in the sequel:

Lemma 3.2. *Let $((\rho_t, m_t^\vartheta) : t \geq 0)$ be a marked exploration process with adapted intensity ϑ.*

i) For each $t > 0$, the increasing càdlàg process $h \mapsto m_t^\vartheta([0, h))$ is $(\mathcal{G}_h^{(t)})_{h\geq 0}$-adapted and its predictable compensator in that filtration is $h \mapsto \int_0^{h\wedge H_t} \vartheta(r, t)\mathrm{d}r$.

ii) For each pair of time instants s and s' one has, almost surely,

$$m_{s'}^\vartheta(\mathrm{d}r)\mathbf{1}_{\{r\leq H_{s,s'}\}} = m_s^\vartheta(\mathrm{d}r)\mathbf{1}_{\{r\leq H_{s,s'}\}}.$$

iii) $((\rho_t, m_t^\vartheta) : t \geq 0)$ is an $(\mathcal{F}_t)$-adapted and càdlàg process with values in $(\mathbb{S}, \hat{d})$.

iv) A marked exploration processes with adapted intensity $((\rho_t, m_t^{\vartheta'}) : t \geq 0)$ is indistinguishable from $((\rho_t, m_t^\vartheta) : t \geq 0)$ if and only if a.s. $\vartheta(t, h) = \vartheta'(t, h)$ a.e. with respect to $\mathrm{d}t \otimes \mathbf{1}_{[0,H_t)}(h)\mathrm{d}h$.

We will refer to property ii) above as *the semi-snake property for marked exploration processes*.

Remark 3.3. Notice that independence of $m_{s'}^\vartheta(\mathrm{d}r)\mathbf{1}_{\{H_{s,s'}<r\leq H_{s'}\}}$ and $m_s^\vartheta(\mathrm{d}r)\mathbf{1}_{\{H_{s,s'}<r\leq H_s\}}$ for $s < s'$ (hence the full "snake-property") will in general fail to hold. Moreover, $((\rho_t, m_t^\vartheta) : t \geq 0)$ is not Markovian in general (an exception is the case of ϑ constant, where the marked exploration process is itself a snake process, with a standard Poisson process as path-space component).

Remark 3.4. Observe that for each $t \geq 0$ the process $h \mapsto \vartheta(h, t)$ is determined from m_t^ϑ only $\mathrm{d}h$ a.e. (as the Lebesgue derivative of its compensator). In view of this and of part iv) of Lemma 3.2, two adapted intensities ϑ and ϑ' are identified if any of the two equivalent properties therein hold.

Proof of Lemma 3.2. Part i) is straightforward using the compensation formula for $\mathcal{N}_t$ with respect to the filtration $(\mathcal{G}_h^{(t)})_{h\geq 0}$, conditionally on $\mathcal{F}_t^\rho = \mathcal{G}_0^{(t)}$.
For part ii), taking conditional expectation given $\mathcal{G}_0^{(s)}$ in the inequality

$$|m_s^\vartheta([0, h)) - m_{s'}^\vartheta([0, h))|\mathbf{1}_{\{h\leq H_{s,s'}\}} \leq \int_0^h \int_{\vartheta(s,r)\wedge\vartheta(s',r)}^{\vartheta(s,r)\vee\vartheta(s',r)} \mathcal{N}_s(\mathrm{d}\nu, \mathrm{d}r)\mathbf{1}_{\{h\leq H_{s,s'}\}},$$

we deduce from $\vartheta(s, \cdot)$ and $\vartheta(s', \cdot)$ being $\mathcal{P}red^{(s)}$-measurable and from the compensation formula with respect to $(\mathcal{G}_t^{(t)})_{r\geq 0}$ that, for each $h \geq 0$, $m_s^\vartheta([0, h)) = m_{s'}^\vartheta([0, h))$ a.s. on $\{h \leq H_{s,s'}\}$. This yields $m_s^\vartheta([0, r)) = m_{s'}^\vartheta([0, r))$ for all rational $r \leq h$ a.s. on $\{h \leq H_{s,s'}\}$, and then $m_s^\vartheta = m_{s'}^\vartheta$ as measures on that event by a.s. left continuity of $r \mapsto m_u^\vartheta([0, r))$, for $u = s, s'$.
Adaptedness in part iii) is clear. As for path regularity of $((\rho_t, m_t^\vartheta) : t \geq 0)$, from the definition (3.2) and the facts that $(\rho_s : s \geq 0)$ is weakly càdlàg and $(H_s : s \geq 0)$ is continuous, we easily get right continuity. Existence of the limit in $(\mathbb{S}, \hat{d})$ of $(\rho_{s_n}, m_{s_n}^\vartheta)$ when $s_n \nearrow t$ is slightly more subtle. Indeed, by path properties of ρ and H, ρ_{s_n} clearly converges to a limit ρ_{t-} such that $H(\rho_{t-}) = H_t$, while

convergence of $m^\vartheta_{s_n}$ to a limit m^ϑ_{t-} follows from completeness of the space of killed càdlàg paths in $\mathbb{R}_+$ under the metric

$$(w,w') \mapsto \int_0^{\zeta(w)\wedge\zeta(w')} \left(\hat{d}_u(w_{(u)}, w'_{(u)}) \wedge 1\right) \mathrm{d}u + \hat{d}_0(w_{(0)}, w'_{(0)}) + |\zeta(w) - \zeta(w')|,$$

where $\zeta(w)$ is the lifetime of w (see [11, Section 4.1.1] for the general property). Last, $\operatorname{supp} m^\vartheta_{t-} \subseteq [0, H(\rho_{t-}))$ follows from the Portmanteau theorem and implies that $(\rho_{t-}, m^\vartheta_{t-}) \in \mathcal{V}$.
For the direct implication in part iv), we first obtain from indistinguishability and part i) the fact that, a.s. for every $t \geq 0$, the processes $h \mapsto \int_0^{h\wedge H_t} \vartheta(r,t)\mathrm{d}r$ and $h \mapsto \int_0^{h\wedge H_t} \vartheta'(r,t)\mathrm{d}r$ are equal. Then, we conclude by Lebesgue derivation. In the converse implication, starting from the inequality

$$|m^\vartheta_s([0,h)) - m^{\vartheta'}_s([0,h))| \leq \int_0^h \int_{\vartheta(s,r)\wedge\vartheta'(s,r)}^{\vartheta(s,r)\vee\vartheta'(s,r)} \mathcal{N}_s(\mathrm{d}\nu, \mathrm{d}r)\,,$$

we obtain (with similar arguments as in proof of point ii) above) the equality $m^\vartheta_s = m^{\vartheta'}_s$ a.s. for all s in a set of full Lebesgue measure. Indistinguishability follows then from the path regularity stated in iii). □

Finally, we establish a useful regularity property regarding the process of marks in each lineage:

Lemma 3.3. *Let $((\rho_t, m^\vartheta_t) : t \geq 0)$ be a marked exploration process with adapted intensity ϑ. Then, the function $s \mapsto m^\vartheta_s([0,H_s))$ is a.s. lower semi-continuous. As a consequence, a.s. the function $s \mapsto \mathbf{1}_{\{m^\vartheta_s = 0\}} = \mathbf{1}_{\{m^\vartheta_s([0,H_s))=0\}}$ has at most countably many discontinuity points.*

Proof. By the semi-snake property of $((\rho_t, m^\vartheta_t) : t \geq 0)$ and the continuity of $s \mapsto H_s$, it holds for each $\varepsilon > 0$ and every $t \geq 0$ that $m^\vartheta_t([0, H_t - \varepsilon)) = m^\vartheta_s([0, H_t - \varepsilon)) \leq m^\vartheta_s([0, H_s))$, for all s close enough to t so that $H_t - \varepsilon \leq H_{s,t}$. Taking $\liminf_{s\to t}$ and letting $\varepsilon \to 0$ gives us the first statement. Also, we deduce that

$$\limsup_{s\to t} \mathbf{1}_{\{m^\vartheta_s([0,H_s))=0\}} \leq \mathbf{1}_{\{m^\vartheta_t([0,H_t))=0\}} \tag{3.3}$$

for all $t \geq 0$, implying that $s \mapsto \mathbf{1}_{\{m^\vartheta_s([0,H_s))=0\}}$ is continuous at points $t \geq 0$ such that $m^\vartheta_t([0,H_t)) > 0$. Now, if $t \geq 0$ is such that $m^\vartheta_t([0,H_t)) = 0$, either we have for any sequence $s_n \to t$ and all large enough n that $\mathbf{1}_{\{m^\vartheta_{s_n}([0,H_{s_n}))=0\}} = 1$, or we can construct a sequence $\hat{s}_n \to t$ such that $\mathbf{1}_{\{m^\vartheta_{\hat{s}_n}([0,H_{\hat{s}_n}))=0\}} = 0$ for all $n \geq 0$. In the first case, $s \mapsto \mathbf{1}_{\{m^\vartheta_s([0,H_s))=0\}}$ is continuous in t. In the second one, t lies in the boundary of the open set $\{s \geq 0 : m^\vartheta_s([0,H_s)) > 0\}$. The latter being a disjoint countable union of open intervals, the conclusion follows. □

3.3 An operator on adapted intensities

The following well known approximations of local time at a given level $a \geq 0$, by the limiting normalized time that the height process spends in a small neighborhood, will be very useful in the sequel:

$$\lim_{\epsilon\to 0} \sup_{a\geq 0} \mathbb{E}\left[\sup_{s\leq t}\left|\epsilon^{-1}\int_0^s \mathbf{1}_{\{a<H_r\leq a+\epsilon\}}dr - L^a_s\right|\right] = 0\,, \tag{3.4}$$

$$\lim_{\epsilon\to 0} \sup_{a\geq \epsilon} \mathbb{E}\left[\sup_{s\leq t}\left|\epsilon^{-1}\int_0^s \mathbf{1}_{\{a-\epsilon<H_r\leq a\}}dr - L^a_s\right|\right] = 0 \tag{3.5}$$

(see [11, Prop. 1.3.3]). We will also need the occupation times formula therefrom deduced or, better, its immediate extension to time-height functions:

$$\int_0^t \varphi(r, H_r)dr = \int_0^\infty \int_0^t \varphi(s,a)dL^a_s da \tag{3.6}$$

a.s. for every $t \geq 0$ and all bounded measurable functions $\varphi : \mathbb{R}_+^2 \to \mathbb{R}$.

Next result will prove the first statement of Theorem 1.2.

Proposition 3.1 (Pruned local times yield adapted intensities). *Let $f : \mathbb{R}_+ \mapsto \mathbb{R}_+$ be a locally bounded measurable function and ϑ an adapted intensity process. Then $(f(L_t^r(m^\vartheta)) : t \geq 0, r \geq 0)$ is also an adapted intensity process.*

Proof. Since $L_t^r(m^\vartheta) \leq L_t^r \leq L_{T_x}^r$ on the event $\{t \leq T_x\}$ and, for each $x \geq 0$ the process $(L_{T_x}^r : r \geq 0)$ is a (sub)critical CSBP issued from x, the fact that $T_x \nearrow \infty$ when $x \to \infty$ implies that a.s. for each $t \geq 0$ the function $r \mapsto L_t^r(m^\vartheta)$ is a.s. locally bounded. This ensures that property 2 in Definition 1.2 is satisfied.

In order to check the first point of Definition 1.2, observe first that for any $\varepsilon, t, r \geq 0$

$$\varepsilon^{-1} \int_0^t \mathbf{1}_{\{-\varepsilon < H_u - r \leq 0\}} \mathrm{d}u = \lim_{k \to \infty} \varepsilon^{-1} \int_0^\infty (1 - \phi_k(u - t + k^{-1}))\, (\phi_k(H_u - r + \varepsilon) - \phi_k(H_u - r))\, \mathrm{d}u\,,$$

almost surely, for $(\phi_k : \mathbb{R} \to [0,1])_{k \geq 1}$ continuous functions vanishing on $(-\infty, 0]$ and equal to 1 on $(1/k, \infty)$. From this and the fact that, for all $t, r \geq 0$ and $k \in \mathbb{N} \backslash \{0\}$, $(H_u \mathbf{1}_{u \leq t - k^{-1}} : u \geq 0)$ is measurable with respect to $\mathcal{B}(\mathbb{R}_+) \otimes \mathcal{G}_0^{(t)} \subset \mathcal{B}(\mathbb{R}_+) \otimes \mathcal{G}_r^{(t)}$, we deduce using (3.5) that $(L_t^r : t \geq 0, r \geq 0)$ is $\mathcal{P}red(\mathcal{N})$-measurable. Since for each $s \geq 0$ the process $r \mapsto m_s^\vartheta([0,r))$ is càglàd and $(\mathcal{G}_r^{(s)})_{r \geq 0}$ adapted, we next observe that

$$(t,r) \mapsto \sum_{k=0}^\infty \left(L^r_{(k+1)2^{-n} \wedge t} - L^r_{k2^{-n} \wedge t} \right) \mathbf{1}_{\{m^\vartheta_{k2^{-n}}([0,r)) = 0\}} = \int_0^t \sum_{k=0}^\infty \mathbf{1}_{\{k2^{-n} < s \leq (k+1)2^{-n},\, m^\vartheta_{k2^{-n}}([0,r)) = 0\}} \mathrm{d}L_s^r$$

is for each $n \in \mathbb{N}$ a $\mathcal{P}red(\mathcal{N})$–measurable process too. Thanks to the a.s. equality $\mathrm{d}L_s^r = \mathbf{1}_{\{H_s = r\}} \mathrm{d}L_s^r$ for each $r \geq 0$, in order to conclude that $(L_t^r(m^\vartheta) : r \geq 0, t \geq 0)$ is $\mathcal{P}red(\mathcal{N})$-measurable it is enough to check the convergence

$$\lim_{n \to \infty} \int_0^t \sum_{k=0}^\infty \mathbf{1}_{\{k2^{-n} < s \leq (k+1)2^{-n}\}} \mathbf{1}_{\{m^\vartheta_{k2^{-n}}([0,r)) = 0\}} \mathrm{d}L_s^r = \int_0^t \mathbf{1}_{\{m_s^\vartheta([0,r)) = 0\}} \mathrm{d}L_s^r \tag{3.7}$$

for each $t \geq 0$. By Lemma 3.3 and continuity of the measure $\mathrm{d}L_s^r$, the function $s \mapsto \mathbf{1}_{\{m_s^\vartheta = 0\}}$ is continuous $\mathrm{d}L_s^r$ almost everywhere. Hence, the functions $s \mapsto \sum_{k=0}^\infty \mathbf{1}_{\{k2^{-n} < s \leq (k+1)2^{-n}\}} \mathbf{1}_{\{m^\vartheta_{k2^{-n}} = 0\}}$ converge $\mathrm{d}L_s^r$ a.e. as $n \to \infty$ to $s \mapsto \mathbf{1}_{\{m_s^\vartheta = 0\}}$. Thus, dominated convergence yields (3.7) and the desired measurability property.

Let us finally check that property 3 in Definition 1.2 holds for $(f(L_t^r(m^\vartheta)) : t \geq 0, r \geq 0)$. Since for all $s' \geq s \geq 0$ and every $r, \varepsilon > 0$ we trivially have $\mathbb{E}\left[\int_s^{s'} \mathbf{1}_{\{r - \varepsilon < H_u \leq r\}} \mathrm{d}u \mathbf{1}_{\{r < H_{s,s'}\}}\right] = 0$, the approximation (3.5) implies that $L_s^r = L_{s'}^r$ for each $r \geq 0$, a.s. on the event $\{r < H_{s,s'}\}$, whence $\int_0^\infty \mathbb{E}(|L_s^r - L_{s'}^r| \mathbf{1}_{\{r < H_{s,s'}\}}) \mathrm{d}r = 0$. Fubini's theorem then ensures that $(L_t^r : t \geq 0, r \geq 0)$ satisfies the semi-snake property for adapted intensities, and the conclusion follows using the approximation (3.7) and the semi-snake property of $(s,r) \mapsto m_s^\vartheta([0,r))$. $\square$

We now establish a continuity-type estimate for the operator F under condition (**H**), that will be useful at several points in the sequel. Introduce for each $M > 0$ the $(\mathcal{F}_t^\rho)_{t \geq 0}$-stopping time

$$T^M := \inf\{t \geq 0 : \sup_{h \geq 0} L_t^h \geq M\} \tag{3.8}$$

and recall that $c(M)$ is a Lipschitz constant of g on $[0, M]$. We have

Lemma 3.4. *Assume condition* (**H**) *holds and let* ϑ_1, ϑ_2 *be two adapted intensity processes. Define for each* $t \geq 0$,

$$\Delta_t := \int_0^{H_t} |\vartheta_1(t,h) - \vartheta_2(t,h)| \mathrm{d}h$$

and

$$\Delta'_t := \int_0^{H_t} |F(\vartheta_1)(t,h) - F(\vartheta_2)(t,h)| \mathrm{d}h \,,$$

Then, for any $M, a \geq 0$ *and every* $(\mathcal{F}^\rho_t)$*-stopping time* τ *such that* $\tau \leq T^M$ *a.s, we have*

$$\mathbb{E}[\mathbf{1}_{\{t\leq\tau,\, H_t\leq a\}}\Delta'_t] \leq c(M)\int_0^t \mathbb{E}[\mathbf{1}_{\{s\leq\tau,\, H_s\leq a\}}\Delta_s]\mathrm{d}s \quad \textit{for all } t \geq 0. \tag{3.9}$$

Proof. Notice first that for each $s \geq 0$, $\left|\mathbf{1}_{\{m^{\vartheta_1}_s=0\}} - \mathbf{1}_{\{m^{\vartheta_2}_s=0\}}\right| \leq \left|m^{\vartheta_1}_s([0,H_s)) - m^{\vartheta_2}_s([0,H_s))\right|$, hence

$$\left|L^h_t(m^{\vartheta_1}) - L^h_t(m^{\vartheta_2})\right| \leq \int_0^t \left|m^{\vartheta_1}_s([0,H_s)) - m^{\vartheta_2}_s([0,H_s))\right|\mathrm{d}L^h_s$$

a.s for all $t \geq 0$. Since for each $h \geq 0$ and for $i = 1, 2$, a.s. $L^h_{t\wedge\tau}(m^{\vartheta_i}) \leq M$ for all $t \geq 0$, we get

$$\begin{aligned}
\mathbf{1}_{\{t\leq\tau,\, H_t\leq a\}}\Delta'_t &\leq c(M)\mathbf{1}_{\{t\leq\tau\}}\int_0^a |L^h_t(m^{\vartheta_1}) - L^h_t(m^{\vartheta_2})|\mathrm{d}h \\
&\leq c(M)\mathbf{1}_{\{t\leq\tau\}}\int_0^a\int_0^t \left|m^{\vartheta_1}_s([0,H_s)) - m^{\vartheta_2}_s([0,H_s))\right|\mathrm{d}L^h_s\mathrm{d}h \\
&\leq c(M)\int_0^a\int_0^t \mathbf{1}_{\{s\leq\tau\}}\left|m^{\vartheta_1}_s([0,h)) - m^{\vartheta_2}_s([0,h))\right|\mathrm{d}L^h_s\mathrm{d}h\Big] \\
&\leq c(M)\int_0^t \mathbf{1}_{\{s\leq\tau,\, H_s\leq a\}}\left|m^{\vartheta_1}_s([0,H_s)) - m^{\vartheta_2}_s([0,H_s))\right|\mathrm{d}s
\end{aligned}$$

using that $\mathrm{d}L^h_s = \mathbf{1}_{\{H_s=h\}}\mathrm{d}L^h_s$ in the third line and the inhomogeneous occupation-time formula (3.6) in the fourth. Thus,

$$\begin{aligned}
\mathbb{E}[\mathbf{1}_{t\leq\tau}\Delta'_t] &\leq c(M)\int_0^t \mathbb{E}\Big[\mathbf{1}_{\{s\leq\tau,\, H_s\leq a\}}\left|m^{\vartheta_1}_s([0,H_s)) - m^{\vartheta_2}_s([0,H_s))\right|\Big]\mathrm{d}s \\
&\leq c(M)\int_0^t \mathbb{E}\left[\mathbf{1}_{\{s\leq\tau,\, H_s\leq a\}}\mathbb{E}\left(\int_0^{H_s}\int_0^\infty |\mathbf{1}_{\{\nu<\vartheta_1(s,h)\}} - \mathbf{1}_{\{\nu<\vartheta_2(s,h)\}}|\mathcal{N}_s(\mathrm{d}h,\mathrm{d}\nu)\middle|\mathcal{F}^\rho_s\right)\right]\mathrm{d}s \\
&\leq c(M)\int_0^t \mathbb{E}\left[\mathbf{1}_{\{s\leq\tau,\, H_s\leq a\}}\mathbb{E}\left(\int_0^{H_s} |\vartheta_1(s,h) - \vartheta_2(s,h)|\mathrm{d}h\middle|\mathcal{F}^\rho_s\right)\right]\mathrm{d}s
\end{aligned}$$

and the statement follows. □

3.4 Iterative scheme and fixed point

The remainder of this section is devoted to the proof of the second part of Theorem 1.2 so from now on we assume that the condition (**H**) holds. The construction of the adapted intensity ϑ^* will be achieved by an iterative procedure. For each $t \geq 0$, we define $\vartheta^0(t,h) = 0$ for all h, so that m^{ϑ^0} is the null measure. Then we let

$$\vartheta^{n+1} = F(\vartheta^n), \forall n \geq 0$$

and write $L^h_t(n) = L^h_t(m^{\vartheta^n})$ to simplify notation.
Observe that, by construction of the sequence ϑ^n, the continuity of g and Lemma 3.1, for all $n \in \mathbb{N}$ the process $(t,h) \mapsto \vartheta^n(t,h)$ has a version which a.s. is continuous in $t \geq 0$ for each $h \geq 0$ and is an adapted intensity process. Moreover, since g is increasing, for any pair $\vartheta, \tilde{\vartheta}$ satisfying a.s. $\vartheta \leq \tilde{\vartheta}$ (pointwise), one has $\mathbf{1}_{\{m^{\vartheta}_t=0\}} \geq \mathbf{1}_{\{m^{\tilde{\vartheta}}_t=0\}}$ for all $t \geq 0$. We can then check by induction in n that a.s. for all $(t,h) \in \mathbb{R}^2_+$,

$$L^h_t(0) \geq L^h_t(2) \geq \cdots \geq L^h_t(2n) \geq \ldots \geq L^h_t(2n+1) \geq \cdots \geq L^h_t(3) \geq L^h_t(1), \tag{3.10}$$

and

$$\vartheta^0(t,h) \leq \vartheta^2(t,h) \leq \cdots \leq \vartheta^{2n}(t,h) \leq \ldots \leq \vartheta^{2n+1}(t,h) \leq \cdots \leq \vartheta^3(t,h) \leq \vartheta^1(t,h). \tag{3.11}$$

Relevant consequences of these inequalities are next highlighted:

Lemma 3.5. *There exist two adapted intensities ϑ^e and ϑ^o and two $\mathcal{P}red(\mathcal{N})$-measurable processes $(L^r_t(e) : t \geq 0, r \geq 0)$, $(L^r_t(o) : t \geq 0, r \geq 0)$ such that, almost surely,*

i)

$$\begin{aligned} \vartheta^{2n} \nearrow \vartheta^e \quad &\text{and} \quad L(2n) \searrow L(e) \\ \vartheta^{2n+1} \searrow \vartheta^o \quad &\text{and} \quad L(2n+1) \nearrow L(o) \end{aligned}$$

pointwise and moreover,

$$\vartheta^e \leq \vartheta^o \text{ and } L(e) \geq L(o)\,,$$

ii) for every $(t,h) \in \mathbb{R}^+_2$ one has $L^h_t(e) = L^h_t(m^{\vartheta^e})$ and $L^h_t(o) = L^h_t(m^{\vartheta^o})$ and

iii) for every $(t,h) \in \mathbb{R}^+_2$ it holds that

$$\begin{cases} \vartheta^e(t,h) = g(L^h_t(m^{\vartheta^o})) \\ \vartheta^o(t,h) = g(L^h_t(m^{\vartheta^e})) \end{cases}.$$

Proof. i) From local boundedness of the process $(t,h) \mapsto L^h_t$ (cf. the proof of Lemma 3.1) and the fact that $\vartheta^n(h,t) \leq g(L^h_t(0))$ for all $n \in \mathbb{N}$ and $(t,h) \in \mathbb{R}^+_2$, the increasing sequence ϑ^{2n} has a pointwise limit ϑ^e. The limit clearly inherits the properties of ϑ^{2n} making it an adapted intensity process. Similar arguments apply to the decreasing sequence ϑ^{2n+1}. Pointwise convergence of the sequences of pruned local times and their measurability properties are also easily obtained by monotone limit arguments. The two inequalities at the end of part i) are immediate consequences of the inequalities (3.10) and (3.11).
We now move to point ii). For fixed $a \geq 0$ and $t \geq 0$, we have

$$L^a_t(e) = \inf_{n\in\mathbb{N}} \int_0^t \mathbf{1}_{\{m^{\vartheta^{2n}}_s([0,H_t))=0\}} \mathrm{d}L^a_s = \int_0^t \inf_{n\in\mathbb{N}} \mathbf{1}_{\{m^{\vartheta^{2n}}_s=0\}} \mathrm{d}L^a_s = \int_0^t \mathbf{1}_{\{m^{\vartheta^e}_s=0\}} \mathrm{d}L^a_s\,,$$

using in the third equality the fact that for each $s \geq 0$ the increasing and integer-valued sequence $m^{\vartheta^{2n}}_s([0,H_s))$ is upper bounded by $m^{\vartheta^1}_s([0,H_s))$, hence constant for large enough n. A similar argument applies for $L^h_t(o)$.
Finally, iii) follows from $\vartheta^e(h,t) = \lim_n \vartheta^{2n}(h,t) = \lim_n F(\vartheta^{2n-1})(h,t) = \lim_n g(L^h_t(2n-1))$. □

Concluding the proof of Theorem 1.2 is now easy:

Proposition 3.2. *With* $\mathbb{P}(d\omega)\otimes \mathrm{d}t\otimes \mathbf{1}_{[0,H_t(\omega))}(h)\mathrm{d}h$ *as reference measure, we have* $\vartheta^e=\vartheta^o=F(\vartheta^e)=F(\vartheta^o)$ *a.e. Moreover,* $\vartheta^*:=\vartheta^e$ *is the a.e. unique solution of the equation* $\vartheta^*=F(\vartheta^*)$ *a.e. and the corresponding marked exploration processes are unique, up to indistinguishability.*

Proof. We apply Lemma 3.4 to $\vartheta_1=\vartheta^e$ and $\vartheta_2=\vartheta^o$. Using Gronwall's lemma and letting M and a therein go to $+\infty$ yields for each $t\geq 0$ that

$$\int_0^{H_t}|\vartheta^e(t,h)-\vartheta^o(t,h)|\mathrm{d}h=0 \quad \text{a.s.}$$

The first statement follows from statement iii) in Lemma 3.5 and integration with respect to $\mathrm{d}t$.
Now, let ϑ^* and ϑ^{**} be two fixed points. By applying Lemma 3.4 to $\vartheta_1=\vartheta^*$ and $\vartheta_2=\vartheta^{**}$, we conclude uniqueness of fixed points using again Gronwall's lemma. Last, point iv) of Lemma 3.2 provides the indistinguishability of the associated marked exploration processes. □

To simplify notation, we will write in the sequel

$$((\rho_t,m_t^*):t\geq 0):=((\rho_t,m_t^{\vartheta^*}):t\geq 0)$$

and refer to this process as the *competitively marked exploration process*. Accordingly, we will call ϑ^* and $L(m^*)$ the *competitive intensity* and *competitively pruned local times*, respectively. Notice that

$$L_t^h(m^*):=\int_0^t \mathbf{1}_{\{m_s^*=0\}}\mathrm{d}L_s^h \tag{3.12}$$

for each $h\geq 0$, a.s. for all $t\geq 0$ and

$$m_t^*([0,h])=\int_0^h\int_0^\infty \mathbf{1}_{\{\nu<g(L_t^r(m^*))\}}\mathcal{N}_t(\mathrm{d}\nu,\mathrm{d}r) \tag{3.13}$$

for each $t\geq 0$, a.s. for all $h\in[0,H_t)$. The rest of the paper is devoted to the proof of Theorem 1.3.

4 A Ray-Knight representation

4.1 Strategy of proof of Theorem 1.3

Our next goal is to prove that the process of $(L_{T_v}^a(m^*):a\geq 0,v\geq 0)$, with $(L_t^a(m^*):t\geq 0,a\geq 0)$ constructed in the previous section and T_x as in (1.10), has the same law as the stochastic flow (1.5).

To that end, we first construct a family $(Z_a^{\varepsilon,\delta}(x):x\geq 0,a\geq 0)$ of approximations of the solution of (1.5), with $(\varepsilon,\delta)\in(0,\infty)^2$, coupled to each other and to the process (1.5) by using the same Gaussian and Poisson noise, and with negative competition drifts that are suitably discretized versions of the drift in (1.5). More precisely, every time instant $a>0$ of the form $k\varepsilon$, $k\in\mathbb{N}$, we split the population into blocks of size δ. For the next $\varepsilon>0$ time units, the evolution of the $n-$th block, $n\in\mathbb{N}$, is given by a flow like (1.3), but with a constant additional negative drift $g(n\delta)$ and suitably "shifted" noises. This construction must be done by means of a nested bi-recursive (in k and n) procedure. The process $(Z_a^{\varepsilon,\delta}(x):x\geq 0,a\geq 0)$ will then be defined by composing the dynamics of consecutive time strips.
Secondly, paralleling the previous construction, we prune $(L_t^a:a\geq 0,t\geq 0)$ at a rate that is constant inside rectangles of some two-dimensional grid, now defined in a bi-recursive way in terms of height and local time units. More precisely, at heights of the form $k\varepsilon$, $k\in\mathbb{N}$, we split the exploration times in blocks of δ units of pruned local time at that level. Then, between height $k\varepsilon$ and $(k+1)\varepsilon$ and when in the $n-$th exploration-time block, we prune the original local time process at rate $g(n\delta)$.

The next step, crucially relying on Proposition 1.2, will be to prove that the resulting punned local times, denoted $(L^a_{T_x}(\varepsilon,\delta) : a \geq 0, x \geq 0)$, has the same law as the process $(Z^{\varepsilon,\delta}_a(x) : a \geq 0, x \geq 0)$. The final and most technical steps will be proving that, when $(\varepsilon,\delta) \to (0,0)$, for each $x \geq 0, a \geq 0$, both the convergences

$$L^a_{T_x}(\varepsilon,\delta) \to L^a_{T_x}(m^*) \quad \text{and} \quad Z^{\varepsilon,\delta}_a(x) \to Z_a(x)$$

hold in probability (in the respective probability spaces). This is enough to grant finite dimensional convergence in distribution and, by the previous step, that $(L^a_{T_x} : a \geq 0, x \geq 0)$ and $(Z_a(x) : x \geq 0, a \geq 0)$ are equal in law as desired.
In the remainder of this section the two approximations are constructed and precise statements on their convergence and on their laws are given (proofs are deferred to Sections 5 and 6). We then end the present section deducing the proof of Theorem 1.3 from those results.

4.2 Grid approximation of the stochastic flow of branching processes with competition

As in (1.5), consider $W(\mathrm{d}s, \mathrm{d}u)$ a white noise process on $(0,\infty)^2$ based on the Lebesgue measure $\mathrm{d}s \otimes \mathrm{d}u$ and N a Poisson random measure on $(0,\infty)^3$ with intensity $\mathrm{d}s \otimes \mathrm{d}\nu \otimes \Pi(\mathrm{d}r)$. Let $\varepsilon, \delta > 0$ be fixed parameters. The process $(Z^{\varepsilon,\delta}_t(x) : x \geq 0, t \geq 0)$ is constructed in the same probability space by means of the following bi-recursive procedure:

Time-step $k = 0$ **:** For all $v \geq 0$, we set

$$Z^{\varepsilon,\delta}_0(v) := Z_0(v) = v;$$

Time-step $k+1$ **:** Assuming that $(Z^{\varepsilon,\delta}_t(w) : t \leq k\varepsilon, w \geq 0)$ is already constructed, we define

$$(Z^{\varepsilon,\delta}_t(w) : t \in (k\varepsilon, (k+1)\varepsilon], w \geq 0)$$

as

$$Z^{\varepsilon,\delta}_t(w) := Z^{\varepsilon,\delta}_{k\varepsilon,t}(Z^{\varepsilon,\delta}_{k\varepsilon}(w)). \tag{4.1}$$

where $(Z^{\varepsilon,\delta}_{k\varepsilon,t}(z) : t \in (k\varepsilon, (k+1)\varepsilon], z \geq 0)$ is an auxiliary process defined as follows:

Space-step $n = 0$ **:** For each $t \in (k\varepsilon, (k+1)\varepsilon]$ we set

$$Z^{\varepsilon,\delta}_{k\varepsilon,t}(0) = 0.$$

Space-step $n+1$ **:** Assuming that a process $(Z^{\varepsilon,\delta}_{k\varepsilon,t}(x) : t \in (k\varepsilon, (k+1)\varepsilon], x \leq n\delta)$ has already been constructed, we set

$$W^{(k,n)}(\mathrm{d}s, \mathrm{d}u) := W(\mathrm{d}s + k\varepsilon, \mathrm{d}\nu + Z_{k\varepsilon,s-}(n\delta))$$

and

$$N^{(k,n)}(\mathrm{d}s, \mathrm{d}\nu, \mathrm{d}r) := N(\mathrm{d}s + k\varepsilon, \mathrm{d}\nu + Z_{k\varepsilon,s-}(n\delta), \mathrm{d}r)$$

which respectively are a Gaussian white noise with intensity $\mathrm{d}s \otimes \mathrm{d}u$ in $\mathbb{R}_+ \times \mathbb{R}_+$ and a Poisson random measure with intensity $\mathrm{d}s \otimes \mathrm{d}\nu \otimes \Pi(\mathrm{d}r)$ in $\mathbb{R}_+ \times \mathbb{R}^2_+$, independent from each other (this can be checked using Lemma 2.1 as in the proof of Proposition 1.1, iii)). We then consider the stochastic flow of $\psi_{g(n\delta)}$-CSBP

$$(Z^{(k,n)}_t(v) : t \geq 0, v \geq 0)$$

defined by

$$\begin{aligned} Z_t^{(k,n)}(v) =& v - \alpha \int_0^t Z_s^{(k,n)}(v)\mathrm{d}s + \sigma \int_0^t \int_0^{Z_{s-}^{(k,n)}(v)} W^{(k,n)}(\mathrm{d}s, \mathrm{d}u) \\ &+ \int_0^t \int_0^{Z_{s-}^{(k,n)}(v)} \int_0^\infty r\tilde{N}^{(k,n)}(\mathrm{d}s, \mathrm{d}\nu, \mathrm{d}r) - \int_0^t g(n\delta) Z_s^{(k,n)}(v)\mathrm{d}s; \end{aligned} \tag{4.2}$$

Here, $\psi_{g(n\delta)}$ is the branching mechanism (1.11) with $\theta = g(n\delta)$, that is,

$$\psi_{g(n\delta)}(\lambda) = \psi(\lambda) + \lambda g(n\delta). \tag{4.3}$$

We then define for all $(t, x) \in (k\varepsilon, (k+1)\varepsilon] \times (n\delta, (n+1)\delta]$:

$$Z_{k\varepsilon,t}^{\varepsilon,\delta}(x) := Z_{k\varepsilon,t}^{\varepsilon,\delta}(n\delta) + Z_{t-k\varepsilon}^{(k,n)}(x - n\delta) = \sum_{l=0}^{n-1} Z_{t-k\varepsilon}^{(k,l)}(l\delta) + Z_{t-k\varepsilon}^{(k,n)}(x - n\delta). \tag{4.4}$$

Remark 4.1. It is immediate that $(W^{(k,n)}(\mathrm{d}s, \mathrm{d}u), N^{(k,n)}(\mathrm{d}s, \mathrm{d}\nu, \mathrm{d}r))_{n\in\mathbb{N}}$ are independent as $k \in \mathbb{N}$ varies. Moreover, reasoning as proof of Proposition 1.1, iii) inductively, one a can also check that for fixed $k \in \mathbb{N}$, $(W^{(k,n)}(\mathrm{d}s, \mathrm{d}u), N^{(k,n)}(\mathrm{d}s, \mathrm{d}\nu, \mathrm{d}r))$ are independent as $n \in \mathbb{N}$ varies. Thus, the processes

$$\left(Z_h^{(k,n)}(v) = Z_{k\varepsilon,k\varepsilon+h}^{\varepsilon,\delta}(n\delta + v) - Z_{k\varepsilon,k\varepsilon+h}^{\varepsilon,\delta}(n\delta) : h \in [0,\varepsilon], v \in [0,\delta] \right)_{k,n\in\mathbb{N}}$$

are independent too.

Definition 4.1. We refer to the processes $(Z_t^{\varepsilon,\delta}(y) : t \geq 0, y \geq 0)$ as the (ε, δ)- grid approximation of the stochastic flow of continuous state branching processes with competition, or simply *grid approximation of the stochastic flow*.

The parameters (ε, δ) being given and fixed, for $x, s \geq 0$ and $k \in \mathbb{N}$ we will use the notation

- $k_s = k_s(\varepsilon) := \sup\{k \in \mathbb{N} : \, k\varepsilon < s\}$ and
- $n_x^k = n_x^k(\varepsilon, \delta) := \sup\{n \in \mathbb{N} : n\delta < Z_{k\varepsilon}^{\varepsilon,\delta}(x)\}$.

By induction in $k \in \mathbb{N}$, summing at each step over $n \in \mathbb{N}$, $(Z_t^{\varepsilon,\delta}(y) : t \geq 0, y \geq 0)$ is seen to solve a sort of stochastic-flow equation, but with a "locally frozen" drift term. Furhtermore, it is not hard to check

Lemma 4.1. *For each $w \geq 0$, the process $Z^{\varepsilon,\delta}(w) = (Z_t^{\varepsilon,\delta}(w) : t \geq 0)$ is a solution to the stochastic differential equation:*

$$\begin{aligned} Z_t^{\varepsilon,\delta}(w) = w - \alpha \int_0^t Z_s^{\varepsilon,\delta}(w)\mathrm{d}s + \sigma \int_0^t \int_0^{Z_{s-}^{\varepsilon,\delta}(w)} W(\mathrm{d}s, \mathrm{d}u) + \int_0^t \int_0^{Z_{s-}^{\varepsilon,\delta}(w)} \int_0^\infty r\tilde{N}(\mathrm{d}s, \mathrm{d}\nu, \mathrm{d}r) \\ - \int_0^t \sum_{n=0}^{n_w^{k_s\varepsilon}} g(n\delta) \left(Z_{k_s\varepsilon,s}((n+1)\delta \wedge Z_{k_s\varepsilon}^{\varepsilon,\delta}(w)) - Z_{k_s\varepsilon,s}(n\delta \wedge Z_{k_s\varepsilon}^{\varepsilon,\delta}(w)) \right), \quad t \geq 0. \end{aligned} \tag{4.5}$$

Moreover, by construction and properties of the flows (4.2) *of CSBP, $(Z_t^{\varepsilon,\delta}(w) : t \geq 0, w \geq 0)$ has a bi-measurable version such that, almost surely, for each $w \geq 0$ the process $t \mapsto Z_t^{\varepsilon,\delta}(w)$ is non-negative and càdlàg and, for every $t \geq 0$, $w \mapsto Z_t^{\varepsilon,\delta}(w)$ is non-negative, non-decreasing and càdlàg.*

The proof of the following result is deferred to Section 5.

Proposition 4.1 (Convergence of the stochastic flow grid approximation). *For each $x \geq 0$ and $t \geq 0$, the r.v. $Z_t^{\varepsilon,\delta}(x)$ converges to $Z_t(x)$ in probability when $(\varepsilon, \delta) \to (0, 0)$.*

4.3 Grid approximation of competitively marked local times

Given fixed parameters $\varepsilon, \delta > 0$, we now introduce an approximation $\vartheta^{\varepsilon,\delta}$ of the intensity process ϑ^*, which will define a pruning at a piecewise constant rate, the regions where it is constant being now the interior of rectangles in a suitably defined height/local-time grid. We will denote by $((\rho_s, m_s^{\varepsilon,\delta}) : s \geq 0)$ the corresponding marked exploration process and by

$$L(\varepsilon, \delta) = (L_t^a(\varepsilon, \delta) : t \geq 0, a \geq 0)$$

the associated pruned local time process, that is, $L_t^a(\varepsilon, \delta) = \int_0^t \mathbf{1}_{\{m_s^{\varepsilon,\delta}=0\}} \mathrm{d}L_s^s$.


ε
$\overline{T}^0_\delta$
$\overline{T}^0_{2\delta}$
2ε
ε
$\overline{T}^\varepsilon_\delta$
$\overline{T}^\varepsilon_{2\delta}$
$\overline{T}^0_\delta$
$\overline{T}^0_{2\delta}$
ε
$\overline{T}^0_\delta$
$\overline{T}^0_{2\delta}$
2ε
ε
$\overline{T}^\varepsilon_\delta$
$\overline{T}^\varepsilon_{2\delta}$
$\overline{T}^0_\delta$
$\overline{T}^0_{2\delta}$

Figure 1: Construction of the local time grid approximation.

Again, the construction consists in a doubly recursive procedure.

Height-step 0 **:** we set $\vartheta^{\varepsilon,\delta}(t, 0) := 0$ for all $t \geq 0$, and $\overline{T}_x^0 = T_x$ for all $x \in \mathbb{R}$.

Height-step $k+1$ **:** Assuming that $\big((\vartheta^{\varepsilon,\delta}(t,h), L_t^h(\varepsilon,\delta)) : t \geq 0, h \in [0, k\varepsilon]\big)$ is constructed, we set for all $x \in \mathbb{R}$,

$$\overline{T}_x^{k\varepsilon} := \inf\{t \geq 0 : L_t^{k\varepsilon}(\varepsilon, \delta) = x\}, \text{ with } \inf \emptyset = \infty.$$

The process $\big((L_t^h(\varepsilon,\delta), \vartheta^{\varepsilon,\delta}(t,h)) : t \geq 0, h \in (k\varepsilon, (k+1)\varepsilon]\big)$ is then defined as follows:

Time-step 0**:** For each $h \in (k\varepsilon, (k+1)\varepsilon]$ we set $\vartheta^{\varepsilon,\delta}(0, h) = g(0)$.

Time-step $n+1$**:** Assume that $\overline{T}_{n\delta}^{k\varepsilon} < \infty$ and that $\Big(\big(L_t^h(\varepsilon,\delta), \vartheta^{\varepsilon,\delta}(t,h)\big) : t \in (0, \overline{T}_{n\delta}^{k\varepsilon}], h \in (k\varepsilon, (k+1)\varepsilon]\big)\Big)$ is already defined. If $\overline{T}_{(n+1)\delta}^{k\varepsilon} < \infty$, for all $(t,h) \in (\overline{T}_{n\delta}^{k\varepsilon}, \overline{T}_{(n+1)\delta}^{k\varepsilon}] \times (k\varepsilon, (k+1)\varepsilon]$ we set

$$\vartheta^{\varepsilon,\delta}(t,h) := g(n\delta), \tag{4.6}$$

$$m_t^{\varepsilon,\delta}([0,h]) := m_t^{\varepsilon,\delta}([0,k\varepsilon]) + \int_{(k\varepsilon,\, h]} \int_0^\infty \mathbf{1}_{\{\nu < g(n\delta)\}} \mathcal{N}_t(\mathrm{d}r, \mathrm{d}\nu), \tag{4.7}$$

and

$$L^h_t(\varepsilon,\delta) := L^h_{\overline{T}^{k\varepsilon}_{n\delta}}(\varepsilon,\delta) + \int_{\overline{T}^{k\varepsilon}_{n\delta}}^t \mathbf{1}_{\{m^{\varepsilon,\delta}_s=0\}}\mathrm{d}L^h_s. \tag{4.8}$$

If $\overline{T}^{k\varepsilon}_{(n+1)\delta} = \infty$, we apply this definitions for all $(t,h) \in (\overline{T}^{k\varepsilon}_{n\delta}, \infty) \times (k\varepsilon, (k+1)\varepsilon]$, and the inductive procedure in n stops. Note that in this case, we have $n\delta \le L^{k\varepsilon}_\infty(\varepsilon,\delta) \le (n+1)\delta$.

It is easily seen that $\vartheta^{\varepsilon,\delta}$ is $\mathcal{P}red(\mathcal{N})$-measurable and that $\vartheta^{\varepsilon,\delta}(s,h) < \vartheta^{\varepsilon,\delta}(s',h)$ can happen for $s \le s'$ only if $L^{k_h\varepsilon}_s < L^{k_h\varepsilon}_{s'}$ and hence $h > H_{s,s'}$. It therefore is an adapted intensity process too.

Remark 4.2. Note that, for each $\varepsilon, \delta > 0$ and $(k,n) \in \mathbb{N}^2$, $L^{k\varepsilon}_{\overline{T}^{k\varepsilon}_{(n+1)\delta}}(\varepsilon,\delta) - L^{k\varepsilon}_{\overline{T}^{k\varepsilon}_{n\delta}}(\varepsilon,\delta)$ takes the value δ if $\overline{T}^{k\varepsilon}_{(n+1)\delta} < \infty$, the value 0 if $\overline{T}^{k\varepsilon}_{n\delta} = \infty$ and some value in $[0,\delta]$ if $\overline{T}^{k\varepsilon}_{n\delta} < \infty$ and $\overline{T}^{k\varepsilon}_{(n+1)\delta} = \infty$.

For fixed parameters (ε,δ) and for $t, h \ge 0$ and $k \in \mathbb{N}$ we introduce the notation

- $\overline{n}^k_t = \overline{n}^k_t(\varepsilon,\delta) := \sup\{n \in \mathbb{N} : n\delta < L^{k\varepsilon}_t(\varepsilon,\delta)\}$,

and recall that we have set $k_h = k_h(\varepsilon) = \sup\{k \in \mathbb{N} : k\varepsilon < h\}$ (see Subsection 4.2). Hence, by construction, a.s for each $t \ge 0$ and all $h \in [0, H_t)$ it holds that

$$\vartheta^{\varepsilon,\delta}(t,h) = g(\overline{n}^{k_h}_t\delta).$$

Moreover, we have

$$L^h_t(\varepsilon,\delta) = \int_0^t \mathbf{1}_{\{m^{\varepsilon,\delta}_s=0\}}\mathrm{d}L^h_s = L^h_t(m^{\varepsilon,\delta}) \tag{4.9}$$

and

$$m^{\varepsilon,\delta}_t([0,h]) = \int_0^h \int_0^\infty \mathbf{1}_{\left\{\nu<\vartheta^{\varepsilon,\delta}(t,r)\right\}}\mathcal{N}_t(\mathrm{d}\nu,\mathrm{d}r) = m^{\vartheta^{\varepsilon,\delta}}_t(h), \tag{4.10}$$

which should be compared to the system of equations (3.12)–(3.13).

Definition 4.2. We refer to the processes $(L^a_t(\varepsilon,\delta) : a \ge 0, t \ge 0)$ as the (ε,δ)- grid approximation of pruned local time or simply *local time grid approximation*.

In what follows, for each fixed height $a \ge 0$ we will denote by $\mathcal{E}_a$ the sigma-field

$$\mathcal{E}_a := \left((\rho_{\tilde{\tau}^a_t}, \mathcal{N}_{\tilde{\tau}^a_t}) : t \ge 0\right) \tag{4.11}$$

where $\tilde{\tau}^a_t$ is the right continuous inverse of $t \mapsto \tilde{A}^a_t := \int_0^t \mathbf{1}_{\{H_s \le a\}}\mathrm{d}s$ (notice that $\mathcal{E}_a$ increases with a).

The following results rely on excursion theory for snake processes and will be proved in Section 6:

Proposition 4.2 (**Law of the local time grid approximation**). *For each fixed $\varepsilon, \delta > 0$ and $(k,n) \in \mathbb{N}^2$, the random variable $L^{k\varepsilon}_{\overline{T}^{k\varepsilon}_{(n+1)\delta}}(\varepsilon,\delta) - L^{k\varepsilon}_{\overline{T}^{k\varepsilon}_{n\delta}}(\varepsilon,\delta)$ is $\mathcal{E}_{k\varepsilon}$-measurable, and the process*

$$\left(L^{k\varepsilon+h}_{\overline{T}^{k\varepsilon}_{n\delta+z}}(\varepsilon,\delta) - L^{k\varepsilon+h}_{\overline{T}^{k\varepsilon}_{n\delta}}(\varepsilon,\delta) : h \in (0,\varepsilon], z \in [0,\delta]\right) \tag{4.12}$$

is $\mathcal{F}_{\overline{T}^{k\varepsilon}_{(n+1)\delta}} \bigwedge \mathcal{E}_{(k+1)\varepsilon}$ measurable. Moreover, the conditional law of the process (4.12) *given $\mathcal{F}_{\overline{T}^{k\varepsilon}_{n\delta}} \bigvee \mathcal{E}_{k\varepsilon}$ is equal to the (unconditional) law of*

$$(Z^{(k,n)}_h(z \wedge y) : h \in [0,\varepsilon], z \in [0,\delta]),$$

when $y = L^{k\varepsilon}_{\overline{T}^{k\varepsilon}_{(n+1)\delta}}(\varepsilon,\delta) - L^{k\varepsilon}_{\overline{T}^{k\varepsilon}_{n\delta}}(\varepsilon,\delta)$ (here, $Z^{(k,n)}$ is a stochastic flow of continuous state branching processes with branching mechanism $\psi_{g(n\delta)}$ as in (4.3)*).*

Lemma 4.2 (Flow property of pruned local-times). *Let ϑ be an adapted intensity and for each $a > 0$ denote by $\breve{T}^a_x := \inf\{t > 0 : \ L^a_t(m^\vartheta) > x\} \leq \infty, x \geq 0$, the right-continuous inverse of the pruned local processes $(L^a_t(m^\vartheta) : t \geq 0)$. Then, for every $a, b, c, x > 0$ we a.s. have*

$$L^{a+b+c}_{\breve{T}^a_x}(m^\vartheta) = L^{a+b+c}_{\breve{T}^{a+b}_{L^{a+b}_{\breve{T}^a_x}(m^\vartheta)}}(m^\vartheta) \quad .$$

Proposition 4.3 (Consistency of the local-time grid approximation). *For each $x \geq 0$ and $a \geq 0$, the random variable $L^a_{T_x}(\varepsilon, \delta)$ converges in probability to $L^a_{T_x}(m^*)$ as $(\varepsilon, \delta) \to 0$.*

4.4 Proof of the Ray-Knight representation

Corollary 4.1. *$(Z^{\epsilon,\delta}_a(x) : x \geq 0, a \geq 0)$ and $(L^a_{T_x}(\epsilon, \delta) : x \geq 0, a \geq 0)$ have the same law.*

Proof. Notice that for every $a, x \geq 0$

$$\begin{aligned} Z^{\varepsilon,\delta}_a(x) &= Z^{\varepsilon,\delta}_{k_a\varepsilon,a}(Z^{\varepsilon,\delta}_{k_a\varepsilon}(x)) = Z^{\varepsilon,\delta}_{k_a\varepsilon,a} \circ Z^{\varepsilon,\delta}_{k_a\varepsilon}(x) \\ &= Z^{\varepsilon,\delta}_{k_a\varepsilon,a} \circ Z^{\varepsilon,\delta}_{(k_a-1)\varepsilon,k_a\varepsilon} \circ Z^{\varepsilon,\delta}_{(k_a-1)\varepsilon}(x) = \cdots \\ &= Z^{\varepsilon,\delta}_{k_a\varepsilon,a} \circ Z^{\varepsilon,\delta}_{(k_a-1)\varepsilon,k_a\varepsilon} \circ \cdots \circ Z^{\varepsilon,\delta}_{0,\varepsilon}(x) \end{aligned}$$

a.s. On the other hand,

$$\begin{aligned} L^a_{T_x}(\varepsilon, \delta) &= L^a_{\overline{T}^{k_a\varepsilon}_{L^{k_a\varepsilon}_{T_x}(\varepsilon,\delta)}}(\varepsilon, \delta) = L^a_{\overline{T}^{k_a\varepsilon}_{\bullet}}(\varepsilon, \delta) \circ L^{k_a\varepsilon}_{T_x}(\varepsilon, \delta) \\ &= L^a_{\overline{T}^{k_a\varepsilon}_{\bullet}}(\varepsilon, \delta) \circ L^{k_a\varepsilon}_{\overline{T}^{(k_a-1)\varepsilon}_{\bullet}}(\varepsilon, \delta) \circ L^{(k_a-1)\varepsilon}_{T_x}(\varepsilon, \delta) = \cdots \\ &= L^a_{\overline{T}^{k_a\varepsilon}_{\bullet}}(\varepsilon, \delta) \circ L^{k_a\varepsilon}_{\overline{T}^{(k_a-1)\varepsilon}_{\bullet}}(\varepsilon, \delta) \circ L^{(k_a-1)\varepsilon}_{\overline{T}^{(k_a-2)\varepsilon}_{\bullet}}(\varepsilon, \delta) \circ \cdots \circ L^{\varepsilon}_{T_x}(\varepsilon, \delta) \end{aligned}$$

for every $a, x \geq 0$ a.s. by Lemma 4.2. It is thus enough to show that, for each $m \in \mathbb{N}$, the equality in law

$$\left(L^{k\varepsilon+h}_{\overline{T}^{k\varepsilon}_y}(\varepsilon, \delta) : h \in [0, \varepsilon], y \geq 0 \right)_{k \leq m} \overset{d}{=} \left(Z^{\varepsilon,\delta}_{k\varepsilon,k\varepsilon+h}(y \wedge Z^{\varepsilon,\delta}_{k\varepsilon}(\infty)) : h \in [0, \varepsilon], y \geq 0 \right)_{k \leq m} \tag{4.13}$$

holds. To that end we proceed by induction on k, with help of the two families of processes

$$\left(A^{(k,n)}(h, z) := L^{k\varepsilon+h}_{\overline{T}^{k\varepsilon}_{n\delta+z}}(\varepsilon, \delta) - L^{k\varepsilon+h}_{\overline{T}^{k\varepsilon}_{n\delta}}(\varepsilon, \delta) : h \in [0, \varepsilon], z \in [0, \delta] \right)_{k,n \in \mathbb{N}}$$

and

$$\left(B^{(k,n)}(h, z) := Z^{(k,n)}_h \left(z \wedge \left(Z^{\epsilon,\delta}_{k\varepsilon}(\infty) - n\delta \right)_+ \right) : h \in [0, \varepsilon], z \in [0, \delta] \right)_{k,n \in \mathbb{N}} .$$

Since $L^0_{\overline{T}^0_{(n+1)\delta}}(\varepsilon, \delta) - L^0_{\overline{T}^0_{n\delta}}(\varepsilon, \delta) = L^0_{T^0_{(n+1)\delta}} - L_{T^0_{n\delta}} = \delta$ a.s. for all $n \in \mathbb{N}$, Proposition 4.2 implies that the processes $\left(A^{(0,n)}\right)_{n \in \mathbb{N}}$ are independent. Thus, $\left(A^{(0,n)}\right)_{n \in \mathbb{N}}$ and $\left(B^{(0,n)}\right)_{n \in \mathbb{N}}$ have the same law by Remark 4.1, and the equality in law (4.13) follows in the case $m = k = 0$ by summing over $n \leq n^0_z$ (note that $Z^{\epsilon,\delta}_0(\infty) = \infty$) for each $x \geq 0$.
We assume next that for $m \geq 1$ the subfamilies $\left(A^{(k,n)}\right)_{k \leq m-1, n \in \mathbb{N}}$ and $\left(B^{(k,n)}\right)_{k \leq m-1, n \in \mathbb{N}}$ have the same law. By summing over n^k_x for each $x \geq 0$ and every $k \leq m-1$, this also grants that the equality in law (4.13) for $m-1$ (instead of m) holds. Let now $(\hat{Z}^{(m,n)})_{n \in \mathbb{N}}$ be a family of processes equal in law to the sequence of processes $(Z^{(m,n)})_{n \in \mathbb{N}}$ defined in (4.2), which are independent by Remark 4.1.

Set $\tilde{n} = \inf\{n \in \mathbb{N} : \overline{T}^{m\varepsilon}_{(n+1)\delta} = \infty\} \leq \infty$ and, for each $m, n \in \mathbb{N}$ and $y \geq 0$, define $\tilde{Z}^{(m,n,y)} = \left(\hat{Z}^{(m,n)}_h(z \wedge y) : h \in [0,\varepsilon], z \in [0,\delta]\right)$. By Proposition 4.2, the conditional law of $\left(A^{(m,n)}\right)_{n\in\mathbb{N}}$ given $\left(A^{(k,l)}\right)_{k\leq m-1, l\in\mathbb{N}}$ is equal to the (unconditional) law of $\left(\tilde{Z}^{(m,n,y_n)}\right)_{n\in\mathbb{N}}$ with $(y_n)_{n\in\mathbb{N}} \in \mathbb{R}_+^{\mathbb{N}}$ given by

$$y_n = \delta \wedge (L^{m\varepsilon}_\infty(\varepsilon,\delta) - n\delta)_+ = \begin{cases} \delta & \text{if } n < \tilde{n} \\ L^{m\varepsilon}_\infty(\delta,\varepsilon) - \tilde{n}\delta & \text{if } n = \tilde{n} \\ 0 & \text{if } n > \tilde{n}. \end{cases}$$

In particular, for every $z \in [0,\delta]$ and $n \in \mathbb{N}$ one has $z \wedge y_n = z \wedge (L^{m\varepsilon}_\infty(\varepsilon,\delta) - n\delta)_+$. The desired equality in law follows then for $k \leq m$ from the induction hypothesis and the fact that

$$L^{m\varepsilon}_\infty(\varepsilon,\delta) = \sum_{n\in\mathbb{N}} A^{(m-1,n)}(\varepsilon,\delta) \quad \text{and} \quad Z^{\varepsilon,\delta}_{m\varepsilon}(\infty) = \sum_{n\in\mathbb{N}} B^{(m-1,n)}(\varepsilon,\delta).$$

□

Proof of Theorem 1.3. By Proposition 4.1 (resp. Proposition 4.3), when $(\varepsilon,\delta) \to (0,0)$ the process $(Z^{\varepsilon,\delta}_a(x) : x \geq 0, a \geq 0)$ (resp. $(L^a_{T_x}(\varepsilon,\delta) : x \geq 0, a \geq 0)$) converges to $(Z_a(x) : x \geq 0, a \geq 0)$ (resp. $(L^a_{T_x}(m^*) : x \geq 0, a \geq 0)$) in the sense of finite dimensional distributions. From these convergences and Corollary 4.1 we conclude that $(Z_a(x) : x \geq 0, a \geq 0)$ and $(L^a_{T_x}(m^*) : x \geq 0, a \geq 0)$ have the same finite dimensional distributions. □

Remark 4.3. Although for all $\varepsilon > 0, \delta > 0$, the process $(Z^{\varepsilon,\delta}_a(x) : x \geq 0, a \geq 0)$ has the same law as $(L^a_{T_x}(\varepsilon,\delta) : x \geq 0, a \geq 0)$ and $(Z_a(x) : x \geq 0, a \geq 0)$ the same as $(L^a_{T_x}(m^*) : x \geq 0, a \geq 0)$, it is not clear whether the couplings $\left((Z^{\varepsilon,\delta}_a(x), Z_a(x)) : x \geq 0, a \geq 0\right)$ and $\left((L^a_{T_x}(\varepsilon,\delta), L^a_{T_x}(m^*)) : x \geq 0, a \geq 0\right)$ are equal in law or not.

5 Consistency of the stochastic flow approximation

5.1 Some auxiliary results

In this section our goal is to prove Proposition 4.1. Recall that the solutions of the various equations (1.3), (1.5) and (4.5) that will be used in its proof are constructed on the same probability space and with the same driving processes. The following comparison property will be useful.

Lemma 5.1 (Comparison property). *Let* $(Y_t(y) : t \geq 0, y \geq 0)$ *be the solution of equation* (1.3). *For all* $\varepsilon, \delta \geq 0$, *the process* $(Z^{\varepsilon,\delta}_t(z) : t \geq 0, z \geq 0)$ *defined by* (4.5) *and the solution* $(Z_t(z) : t \geq 0, z \geq 0)$ *to equation* (1.5) *satisfy*

$$\mathbb{P}\{Z^{\varepsilon,\delta}_t(v) \leq Y_t(w), \forall t \geq 0\} = 1 \quad \textit{and} \quad \mathbb{P}\{Z_t(v) \leq Y_t(w), \forall t \geq 0\} = 1, \tag{5.1}$$

for all $0 \leq v \leq w$. *In both cases we say that the "comparison property" holds.*

Proof. The comparison property for $Z(v)$ with respect to $Y(w)$ follows directly from [10, Theorem 2.2]. Whenever $y \geq z$, the same result implies a comparison property for each of the $\psi_{g(n\delta)}$-CSBP $(Z^{(k,n)}_t(z) : t \geq 0)$ defined in (4.2), with respect to a ψ-CSBP $(Y^{(k,n)}_t(y) : t \geq 0)$ constructed using *the same noises* $(W^{(k,n)}(\mathrm{d}s, \mathrm{d}u), N^{(k,n)}(\mathrm{d}s, \mathrm{d}\nu, \mathrm{d}r))_{n\in\mathbb{N}}$. Since $\left(Z^{\varepsilon,\delta}_{k\varepsilon,k\varepsilon+h}(z) : h \in [0,\varepsilon]\right)$ is build for each k by summing processes $(Z^{(k,n)}_t(x) : t \geq 0)$ defined in (4.4), over finitely many n and for suitable values of x, an inductive argument in n yields the comparison property of that process with respect to the process $\big(Y_{k\varepsilon,k\varepsilon+h}(y) : h \in [0,\varepsilon]\big)$ defined in (1.4), which can be obtained by similarly summing over n some of the above described processes $(Y_t^{(k,n)}(x) : t \geq 0)$. Last, $(Z_t^{\varepsilon,\delta}(v) : t \geq 0, v \geq 0)$ and $(Y_t(w) : t \geq 0, w \geq 0)$ can be respectively obtained from the families $(Z^{\varepsilon,\delta}_{k\varepsilon,k\varepsilon+\cdot})_{k\in\mathbb{N}}$ and $(Y_{k\varepsilon,k\varepsilon+\cdot})_{k\in\mathbb{N}}$ by composition, so an induction argument in k completes the proof of the desired property. $\square$

To unburden the proof of Proposition 4.1, we next prepare a technical lemma.

Lemma 5.2. *Let $x \geq 0$ be fixed and for $m \geq 0$ let $\tau_m = \tau_m(x) := \inf\{t \geq 0 : Y_t(x) > m\}$. Then, for all $s \geq 0$,*

$$\mathbb{E}\left[\mathbf{1}_{\{s\leq\tau_m\}} \sup_{0\leq v\leq Z^{\varepsilon,\delta}_{k_s\varepsilon}(x)} \left|Z^{\varepsilon,\delta}_{k_s\varepsilon,s}(v) - v\right|\right] \leq m\varepsilon\left(|\alpha_0| + \int_1^\infty r\Pi(\mathrm{d}r)\right) + g(m)m\varepsilon \\ + \sqrt{m\varepsilon}C_1\left(\sigma + \sqrt{\int_0^1 r^2\Pi(\mathrm{d}r)}\right),$$

where $\alpha_0 = -\alpha - \int_1^\infty x\Pi(dx)$ and $C_1 > 0$ is some universal constant.

Proof. It follows from (4.2) and (4.4) that

$$\begin{aligned}
\mathbb{E}\left[\mathbf{1}_{\{s\leq\tau_m\}} \sup_{0\leq v\leq Z^{\varepsilon,\delta}_{k_s\varepsilon}(x)} \left|Z^{\varepsilon,\delta}_{k_s\varepsilon,s}(v) - v\right|\right] \leq& \mathbb{E}\left[\mathbf{1}_{\{s\leq\tau_m\}}\left(|\alpha_0|\int_{k_s\varepsilon}^s Z^{\varepsilon,\delta}_\theta(x)\mathrm{d}\theta + \int_{k_s\varepsilon}^s\int_0^{Z^{\varepsilon,\delta}_\theta(x)}\int_1^\infty rN(\mathrm{d}\theta,\mathrm{d}\nu,\mathrm{d}r)\right)\right] \\
&+ \sigma\mathbb{E}\left[\mathbf{1}_{\{s\leq\tau_m\}} \sup_{0\leq v\leq Z^{\varepsilon,\delta}_{k_s\varepsilon}(x)} \left|\int_{k_s\varepsilon}^s\int_0^{Z^{\varepsilon,\delta}_{k_s\varepsilon,\theta}(v)} W(\mathrm{d}\theta,\mathrm{d}u)\right|\right] \\
&+ \mathbb{E}\left[\mathbf{1}_{\{s\leq\tau_m\}} \sup_{0\leq v\leq Z^{\varepsilon,\delta}_{k_s\varepsilon}(x)} \left|\int_{k_s\varepsilon}^s\int_0^{Z^{\varepsilon,\delta}_{k_s\varepsilon,\theta}(v)}\int_0^1 r\tilde{N}(\mathrm{d}\theta,\mathrm{d}\nu,\mathrm{d}r)\right|\right] \\
&+ \mathbb{E}\left[\mathbf{1}_{\{s\leq\tau_m\}} g(Z^{\varepsilon,\delta}_{k_s\varepsilon}(x))\int_{k_s\varepsilon}^s Z^{\varepsilon,\delta}_\theta(x)\mathrm{d}\theta\right].
\end{aligned}$$

Thanks to Lemma 5.1, on $\{s \leq \tau_m\}$ it holds a.s. for every $(t,v) \in (k_s\varepsilon, s]\times[0, Z^{\varepsilon,\delta}_{k_s\varepsilon}(x))$ that $Z^{\varepsilon,\delta}_{k_s\varepsilon,t}(v) = Z^{\varepsilon,\delta}_{k_s\varepsilon,t}(v) \wedge m$. We deduce

$$\begin{aligned}
\mathbb{E}\left[\mathbf{1}_{\{s\leq\tau_m\}} \sup_{0\leq v\leq Z^{\varepsilon,\delta}_{k_s\varepsilon}(x)} \left|Z^{\varepsilon,\delta}_{k_s\varepsilon,s}(v) - v\right|\right] \leq& \sigma\mathbb{E}\left[\sup_{0\leq v\leq Z^{\varepsilon,\delta}_{k_s\varepsilon}(x)\wedge m} |M_v^W(s)|\right] + \mathbb{E}\left[\sup_{0\leq v\leq Z^{\varepsilon,\delta}_{k_s\varepsilon}(x)\wedge m} |M_v^N(s)|\right] \\
&+ m\varepsilon\left(|\alpha_0| + \int_1^\infty r\Pi(\mathrm{d}r)\right) + g(m)m\varepsilon
\end{aligned} \tag{5.2}$$

where, for each $s \geq 0$, the processes $(M_v^W(s))_{v\geq0}$ and $(M_v^N(s))_{v\geq0}$ are respectively defined as

$$M_v^W(s) = \int_{k_s\varepsilon}^s\int_0^{Z^{\varepsilon,\delta}_{k_s\varepsilon,\theta}(v)\wedge m} W(\mathrm{d}\theta,\mathrm{d}u) \quad \text{and} \quad M_v^N(s) = \int_{k_s\varepsilon}^s\int_0^{Z^{\varepsilon,\delta}_{k_s\varepsilon,\theta}(v)\wedge m}\int_0^1 r\tilde{N}(\mathrm{d}\theta,\mathrm{d}\nu,\mathrm{d}r).$$

The two of them are martingales in the variable v started from 0, with respect to the filtration $(\mathcal{S}_v^{k_s\varepsilon,s})_{v\geq 0}$ given by

$$\mathcal{S}_v^{k_s\varepsilon,s} := \sigma\left(\mathbf{1}_{[0,Z^{\varepsilon,\delta}_{k_s\varepsilon,t}(w)]}(u)W(\mathrm{d}\theta,\mathrm{d}u), \mathbf{1}_{[0,Z^{\varepsilon,\delta}_{k_s\varepsilon,t}(w)]}(\nu)N(\mathrm{d}\theta,\mathrm{d}\nu,\mathrm{d}r) : \theta\in[k_s\varepsilon,s], 0\leq w\leq v\right).$$

Indeed, we have

$$\begin{aligned}
&\mathbb{E}\left[\int_{k_s\varepsilon}^{s}\int_0^{Z^{\varepsilon,\delta}_{k_s\varepsilon,\theta}(v+h)\wedge m} W(\mathrm{d}\theta,\mathrm{d}u)\middle|\mathcal{S}_v^{k_s\varepsilon,s}\right] \\
&\qquad= \mathbb{E}\left[\int_{k_s\varepsilon}^{s}\int_0^{Z^{\varepsilon,\delta}_{k_s\varepsilon,\theta}(v)\wedge m} W(\mathrm{d}\theta,\mathrm{d}u)\middle|\mathcal{S}_v^{k_s\varepsilon,s}\right] + \mathbb{E}\left[\int_{k_s\varepsilon}^{s}\int_{Z^{\varepsilon,\delta}_{k_s\varepsilon,\theta}(v)\wedge m}^{Z^{\varepsilon,\delta}_{k_s\varepsilon,\theta}(v+h)\wedge m} W(\mathrm{d}\theta,\mathrm{d}u)\middle|\mathcal{S}_v^{k_s\varepsilon,s}\right] \\
&\qquad= \int_{k_s\varepsilon}^{s}\int_0^{Z^{\varepsilon,\delta}_{k_s\varepsilon,\theta}(v)\wedge m} W(\mathrm{d}\theta,\mathrm{d}u) + \mathbb{E}\left[\int_{k_s\varepsilon}^{s}\int_{Z^{\varepsilon,\delta}_{k_s\varepsilon,\theta}(v)\wedge m}^{Z_{k_s\varepsilon,\theta}(v+h)\wedge m} W(\mathrm{d}\theta,\mathrm{d}u)\right] \\
&\qquad= \int_{k_s\varepsilon}^{s}\int_0^{Z^{\varepsilon,\delta}_{k_s\varepsilon,\theta}(v)\wedge m} W(\mathrm{d}\theta,\mathrm{d}u),
\end{aligned}$$

where the second equality can be checked using Remark 4.1 and the flow property of the $\psi_{g(n\delta)}$-CSBPs (4.2). In a similar way, we get that

$$\mathbb{E}\left[\int_{k_s\varepsilon}^{s}\int_0^{Z^{\varepsilon,\delta}_{k_s\varepsilon,\theta}(v+h)\wedge m}\int_0^1 r\tilde{N}(\mathrm{d}\theta,\mathrm{d}\nu,\mathrm{d}r)\middle|\mathcal{S}_v^{k_s\varepsilon,s}\right] = \int_{k_s\varepsilon}^{s}\int_0^{Z^{\varepsilon,\delta}_{k_s\varepsilon,\theta}(v)\wedge m}\int_0^1 r\tilde{N}(\mathrm{d}\theta,\mathrm{d}\nu,\mathrm{d}r).$$

By Ito's formula in the time variable $s\geq 0$, we then see that

$$\begin{aligned}
\mathbb{E}\left[M_v^W(s)^2\right] &= 2\mathbb{E}\left[\int_{k_s\varepsilon}^{s}\int_0^{Z^{\varepsilon,\delta}_{k_s\varepsilon,\theta}(v)\wedge m} M_u^W(\theta)W(\mathrm{d}\theta,\mathrm{d}u)\right] + \mathbb{E}\left[\int_{k_s\varepsilon}^{s}(Z^{\varepsilon,\delta}_{k_s\varepsilon,\theta}(v)\wedge m)\mathrm{d}\theta\right] \\
&= \mathbb{E}\left[\int_{k_s\varepsilon}^{s}(Z^{\varepsilon,\delta}_{k_s\varepsilon,\theta}(v)\wedge m)\mathrm{d}\theta\right],
\end{aligned}$$

which implies that

$$\mathbb{E}\left[\left[M_\cdot^W(s), M_\cdot^W(s)\right]_v\right] = \mathbb{E}\left[\int_{k_s\varepsilon}^{s}(Z^{\varepsilon,\delta}_{k_s\varepsilon,\theta}(v)\wedge m)\mathrm{d}\theta\right]$$

by definition of the quadratic variation of $(M_v^W(s))_{v\geq 0}$. Thus, conditioning first on the sigma field generated by W and N up to time $k_s\varepsilon$ and using then Burkholder-Davis-Gundy's inequality in the spatial variable $v\geq 0$, the first term on the right hand side of (5.2) can be bounded as follows:

$$\begin{aligned}
\sigma\mathbb{E}\left[\sup_{0\leq v\leq Z^{\varepsilon,\delta}_{k_s\varepsilon}(x)\wedge m}|M_v^W(s)|\right] &\leq \sigma C_1\mathbb{E}\left[\sqrt{[M_\cdot^W(s), M_\cdot^W(s)]_{(Z^{\varepsilon,\delta}_{k_s\varepsilon}(x)\wedge m)}}\right] \\
&\leq \sigma C_1\sqrt{\mathbb{E}\left[\int_{k_s\varepsilon}^{s}(Z^{\varepsilon,\delta}_\theta(x)\wedge m)\mathrm{d}\theta\right]} \leq \sigma C_1\sqrt{m\varepsilon},
\end{aligned}\tag{5.3}$$

with $C_1>0$ a universal constant. Similarly, by Itô's formula,

$$\begin{aligned}
\mathbb{E}\left[M_v^{\tilde{N}}(s)^2\right] =& 2\mathbb{E}\left[\int_{k_s\varepsilon}^{s}\int_0^{Z_{k_s\varepsilon,\theta}(v)\wedge m}\int_0^1 M_u^{\tilde{N}}(\theta)\tilde{N}(\mathrm{d}\theta,\mathrm{d}\nu,\mathrm{d}r)\right] + \mathbb{E}\left[\sum_{k_s\varepsilon<s_n\leq s} r_n^2\mathbf{1}_{\{\nu_n\leq(Z_{k_s\varepsilon,s_n}(v)\wedge m)\}}\right] \\
=& \mathbb{E}\left[\int_{k_s\varepsilon}^{s}\int_0^{Z_{k_s\varepsilon,\theta}(v)\wedge m}\int_0^1 r^2\Pi(\mathrm{d}r)\mathrm{d}\theta\mathrm{d}\nu\right] = \mathbb{E}\left[\left[M_\cdot^{\tilde{N}}(s), M_\cdot^{\tilde{N}}(s)\right]_v\right].
\end{aligned}$$

In a similar way as before, for the second term on the right hand side of (5.2) we get the bound

$$
\begin{aligned}
\mathbb{E}\left[\sup_{0\le v\le Z^{\varepsilon,\delta}_{k_s\varepsilon}(x)\wedge m}|M^{\tilde{N}}_v(s)|\right] &\le C_1\mathbb{E}\left[\sqrt{\left[M^{\tilde{N}}_\cdot(s), M^{\tilde{N}}_\cdot(s)\right]_{Z^{\varepsilon,\delta}_{k_s\varepsilon}(x)\wedge m}}\right] \\
&\le C_1\sqrt{\mathbb{E}\left[\int_{k_s\varepsilon}^{s}(Z^{\varepsilon,\delta}_\theta(x)\wedge m)\mathrm{d}\theta\int_0^1 r^2\Pi(\mathrm{d}r)\right]} \\
&\le \sqrt{m\varepsilon}C_1\sqrt{\int_0^1 r^2\Pi(\mathrm{d}r)}.
\end{aligned}
\tag{5.4}
$$

Using the bounds (5.3) and (5.4) in (5.2) the desired result follows. □

5.2 Proof of Proposition 4.1

Proof of Proposition 4.1. The difference $\zeta^{\varepsilon,\delta}_t(x) := Z_t(x) - Z^{\varepsilon,\delta}_t(x)$ satisfies

$$
\begin{aligned}
\zeta^{\varepsilon,\delta}_t(x) = &-\alpha\int_0^t\left(Z_s(x)-Z^{\varepsilon,\delta}_s(x)\right)\mathrm{d}s + \sigma\int_0^t\int_0^\infty\left(\mathbf{1}_{\{u<Z_{s-}(x)\}}-\mathbf{1}_{\{u<Z^{\varepsilon,\delta}_{s-}(x)\}}\right)W(\mathrm{d}s,\mathrm{d}u) \\
&+\int_0^t\int_0^\infty\int_0^\infty\left(\mathbf{1}_{\{\nu<Z_{s-}(x)\}}-\mathbf{1}_{\{\nu<Z^{\varepsilon,\delta}_{s-}(x)\}}\right)r\tilde{N}(\mathrm{d}s,\mathrm{d}\nu,\mathrm{d}r) \\
&-\int_0^t\left[G(Z_s(x))-G(Z^{\varepsilon,\delta}_s(x))\right]\mathrm{d}s-\int_0^t\int_0^{Z^{\varepsilon,\delta}_s(x)}g(v)\mathrm{d}v\mathrm{d}s \\
&+\int_0^t\sum_{n=0}^{n_x^{k_s\varepsilon}}g(n\delta)\left(Z_{k_s\varepsilon,s}((n+1)\delta\wedge Z^{\varepsilon,\delta}_{k_s\varepsilon}(x))-Z_{k_s\varepsilon,s}(n\delta\wedge Z^{\varepsilon,\delta}_{k_s\varepsilon}(x))\right)\mathrm{d}s.
\end{aligned}
\tag{5.5}
$$

Observe that

$$
Z_{k_s\varepsilon,s}((n+1)\delta\wedge Z^{\varepsilon,\delta}_{k_s\varepsilon}(x))-Z_{k_s\varepsilon,s}(n\delta\wedge Z^{\varepsilon,\delta}_{k_s\varepsilon}(x)) = \int_{n\delta\wedge Z^{\varepsilon,\delta}_{k_s\varepsilon}(x)}^{(n+1)\delta\wedge Z^{\varepsilon,\delta}_{k_s\varepsilon}(x)}\mathrm{d}_u Z_{k_s\varepsilon,s}(u),
$$

which we can substitute in the last term in the right-hand side of (5.5) to get

$$
\begin{aligned}
\zeta^{\varepsilon,\delta}_t(x) = &-\alpha\int_0^t\left(Z_s(x)-Z^{\varepsilon,\delta}_s(x)\right)\mathrm{d}s + \sigma\int_0^t\int_0^\infty\left(\mathbf{1}_{\{u<Z_{s-}(x)\}}-\mathbf{1}_{\{u<Z^{\varepsilon,\delta}_{s-}(x)\}}\right)W(\mathrm{d}s,\mathrm{d}u) \\
&+\int_0^t\int_0^\infty\int_0^\infty\left(\mathbf{1}_{\{\nu<Z_{s-}(x)\}}-\mathbf{1}_{\{\nu<Z^{\varepsilon,\delta}_{s-}(x)\}}\right)r\tilde{N}(\mathrm{d}s,\mathrm{d}\nu,\mathrm{d}r) \\
&-\int_0^t\left[G(Z_s(x))-G(Z^{\varepsilon,\delta}_s(x))\right]\mathrm{d}s \\
&-\int_0^t\sum_{n=0}^{n_x^{k_s\varepsilon}}\int_0^{Z^{\varepsilon,\delta}_{k_s\varepsilon}(x)}\mathbf{1}_{\{n\delta<u\le(n+1)\delta\}}\left[g(u)-g(n\delta)\right]d_u\,Z^{\varepsilon,\delta}_{k_s\varepsilon,s}(u)\mathrm{d}s \\
&-\int_0^t\left[\int_0^{Z^{\varepsilon,\delta}_s(x)}g(v)\mathrm{d}v-\int_0^{Z^{\varepsilon,\delta}_{k_s\varepsilon}(x)}g(u)\mathrm{d}_u Z^{\varepsilon,\delta}_{k_s\varepsilon,s}(u)\right]\mathrm{d}s.
\end{aligned}
\tag{5.6}
$$

Applying Itô's formula to an approximation of the absolute value as in the proof of [10, Theorem 2.1] and dealing with second order terms (5.6) in a similar way, we deduce (see Appendix A.2 for details) that for each $m \in \mathbb{N}$,

$$
\begin{aligned}
\mathbb{E}|\zeta^{\varepsilon,\delta}_{t\wedge\tau_m}(x)| \le \Big(|\alpha_0| + \int_1^\infty r\Pi(dr) + g(m)\Big)\, \mathbb{E}\left[\int_0^t |\zeta^{\varepsilon,\delta}_{s\wedge\tau_m}(x)|\mathrm{d}s\right] \\
+ \mathbb{E}\left[\int_0^{t\wedge\tau_m} \sum_{n=0}^{n_x^{k_s\varepsilon}} \int_0^{Z^{\varepsilon,\delta}_{k_s\varepsilon}(x)} \mathbf{1}_{\{n\delta<u\le(n+1)\delta\}}\, |g(u)-g(n\delta)|\, d_u Z^{\varepsilon,\delta}_{k_s\varepsilon,s}(u)\mathrm{d}s\right] \\
+ \mathbb{E}\int_0^{t\wedge\tau_m}\left|\int_0^{Z^{\varepsilon,\delta}_s(x)} g(v)\mathrm{d}v - \int_0^{Z^{\varepsilon,\delta}_{k_s\varepsilon}(x)} g(v)\mathrm{d}v\right|\mathrm{d}s \\
+ \mathbb{E}\int_0^{t\wedge\tau_m}\left|\int_0^{Z^{\varepsilon,\delta}_{k_s\varepsilon}(x)} g(u)\mathrm{d}u - \int_0^{Z^{\varepsilon,\delta}_{k_s\varepsilon}(x)} g(u)\mathrm{d}_u Z^{\varepsilon,\delta}_{k_s\varepsilon,s}(u)\right|\mathrm{d}s.
\end{aligned}
\tag{5.7}
$$

Since $|u-n\delta|\,\mathbf{1}_{\{n\delta<u\le(n+1)\delta\}} \le \delta$ and $Z^{\varepsilon,\delta}_{k_s\varepsilon,s}(Z^{\varepsilon,\delta}_{k_s\varepsilon}(x)) = Z^{\varepsilon,\delta}_s(x) \le m$ for $s \le \tau_m$, the second term on the right hand side of (5.7) is bounded by $c(m)m\delta t$. We can use integration by parts inside the last term of (5.7) in order to rewrite

$$
\int_0^{Z^{\varepsilon,\delta}_{k_s\varepsilon}(x)} g(u)\mathrm{d}u = g(Z^{\varepsilon,\delta}_{k_s\varepsilon}(x))Z^{\varepsilon,\delta}_{k_s\varepsilon}(x) - \int_0^{Z^{\varepsilon,\delta}_{k_s\varepsilon}(x)} u\, g'(u)\mathrm{d}u
$$

and

$$
\begin{aligned}
-\int_0^{Z^{\varepsilon,\delta}_{k_s\varepsilon}(x)} g(u)\mathrm{d}_u Z^{\varepsilon,\delta}_{k_s\varepsilon,s}(u) &= -g(u)Z^{\varepsilon,\delta}_{k_s\varepsilon,s}(u)\Big|_{u=0}^{u=Z^{\varepsilon,\delta}_{k_s\varepsilon}(x)} + \int_0^{Z^{\varepsilon,\delta}_{k_s\varepsilon}(x)} Z^{\varepsilon,\delta}_{k_s\varepsilon,s}(u)g'(u)\mathrm{d}u \\
&= -g(Z^{\varepsilon,\delta}_{k_s\varepsilon}(x))Z^{\varepsilon,\delta}_s(x) + \int_0^{Z^{\varepsilon,\delta}_{k_s\varepsilon}(x)} Z^{\varepsilon,\delta}_{k_s\varepsilon,s}(u)g'(u)\mathrm{d}u
\end{aligned}
$$

(since $u \mapsto g(u)$ is locally Lipschitz), and then deduce the following upper bound for the sum of the third and fourth lines therein:

$$
\begin{aligned}
\mathbb{E}\int_0^{t\wedge\tau_m}\Big(g(m) + g(Z^{\varepsilon,\delta}_{k_s\varepsilon}(x))\Big)\left|Z^{\varepsilon,\delta}_{k_s\varepsilon}(x) - Z^{\varepsilon,\delta}_s(x)\right|\mathrm{d}s \\
+ \mathbb{E}\int_0^{t\wedge\tau_m}\int_0^{Z^{\varepsilon,\delta}_{k_s\varepsilon}(x)}\left|u - Z^{\varepsilon,\delta}_{k_s\varepsilon,s}(u)\right| g'(u)\mathrm{d}u\mathrm{d}s.
\end{aligned}
$$

Summarizing, we have

$$
\begin{aligned}
\mathbb{E}|\zeta^{\varepsilon,\delta}_{t\wedge\tau_m}(x)| \le \Big(|\alpha_0| + \int_1^\infty r\Pi(\mathrm{d}r) + g(m)\Big)\mathbb{E}\left[\int_0^t |\zeta^{\varepsilon,\delta}_{s\wedge\tau_m}(x)|\mathrm{d}s\right] + c(m)m\delta t \\
+\mathbb{E}\int_0^{t\wedge\tau_m}\Big(g(m) + g(Z^{\varepsilon,\delta}_{k_s\varepsilon}(x))\Big)\left|Z^{\varepsilon,\delta}_{k_s\varepsilon}(x) - Z^{\varepsilon,\delta}_s(x)\right|\mathrm{d}s \\
+\mathbb{E}\int_0^{t\wedge\tau_m}\int_0^{Z^{\varepsilon,\delta}_{k_s\varepsilon}(x)}\left|u - Z^{\varepsilon,\delta}_{k_s\varepsilon,s}(u)\right| g'(u)\mathrm{d}u\mathrm{d}s \\
\le \Big(|\alpha_0| + \int_1^\infty r\Pi(\mathrm{d}r) + g(m)\Big)\mathbb{E}\left[\int_0^t |\zeta^{\varepsilon,\delta}_{s\wedge\tau_m}(x)|\mathrm{d}s\right] + c(m)m\delta t \\
+2g(m)\mathbb{E}\int_0^{t\wedge\tau_m}\left|Z^{\varepsilon,\delta}_{k_s\varepsilon}(x) - Z^{\varepsilon,\delta}_{k_s\varepsilon,s}(Z^{\varepsilon,\delta}_{k_s\varepsilon}(x))\right|\mathrm{d}s \\
+c(m)\mathbb{E}\int_0^{t\wedge\tau_m}\int_0^{Z^{\varepsilon,\delta}_{k_s\varepsilon}(x)}\left|u - Z^{\varepsilon,\delta}_{k_s\varepsilon,s}(u)\right|\mathrm{d}u\mathrm{d}s.
\end{aligned}
\tag{5.8}
$$

We then use Lemma 5.2 to bound the last two terms in (5.8) and get that

$$\begin{aligned}\mathbb{E}|\zeta^{\varepsilon,\delta}_{t\wedge\tau_m}(x)| \leq & \left(|\alpha_0| + \int_1^\infty r\Pi(\mathrm{d}r) + g(m)\right)\mathbb{E}\left[\int_0^t |\zeta^{\varepsilon,\delta}_{s\wedge\tau_m}(x)|\mathrm{d}s\right] + c(m)m\delta t \\ & + (g(m)+2c(m))\, tm\varepsilon\left(|\alpha_0| + \int_1^\infty r\Pi(\mathrm{d}r) + c(m)\right) \\ & + (g(m)+2c(m))\, t\sqrt{m\varepsilon}C_1\left(\sigma + \sqrt{\int_0^1 r^2\Pi(\mathrm{d}r)}\right)\end{aligned}$$

Since $\zeta^{\varepsilon,\delta}_s(x) \leq 2m$ when $0 < s \leq \tau_m$ by Lemma 5.1, we can use Gronwall's lemma and get

$$\begin{aligned}\mathbb{E}|\zeta^{\epsilon,\delta}_{t\wedge\tau_m}(x)| \leq & \Bigg[\delta c(m)m + \varepsilon\,(g(m)+2c(m))\, m\left(|\alpha_0| + \int_1^\infty r\Pi(\mathrm{d}r) + c(m)\right) \\ & + \sqrt{\varepsilon}\,(g(m)+2c(m))\sqrt{m}C_1\left(\sigma + \sqrt{\int_0^1 r^2\Pi(\mathrm{d}r)}\right)\Bigg]\, te^{(|\alpha_0|+\int_1^\infty r\Pi(\mathrm{d}r)+g(m))t}.\end{aligned}$$

Hence, $\mathbb{E}|\zeta^{\epsilon,\delta}_{t\wedge\tau_m}(x)|$ goes to zero when $(\delta,\varepsilon)\to(0,0)$ for each fixed $m>0$. The desired result follows since $\tau_m\to\infty$ a.s. as $m\to\infty$ □

6 Law and consistency of the local time approximation

The proofs of Propositions 4.2 and 4.3 are based on several technical results mainly relying on excursion theory and on Proposition 1.2. For easier reading, some proofs are differed to the Appendix.
We will need in what follows the $(\mathcal{F}^\rho_t)$-stopping times defined by

$$T^a_x := \inf\{t \geq 0 : L^a_t \geq x\}$$

so that $T^0_x = T_x$ for all $x \geq 0$.

6.1 Poisson-snake excursions

We first briefly recall some facts of excursion theory in the case of the snake process $((\rho_t,\mathcal{N}_t) : t\geq 0)$. The reader is referred to [11, 12] for details and general background.
Recall that the processes $(X_s - I_s : s\geq 0) = (\langle\rho_s, 1\rangle : s \geq 0)$ and ρ share the same excursion intervals $(\alpha_j,\beta_j)_{j\in J}$ away from their respective zero elements. Let us denote by $(\rho^j,\mathcal{N}^j)$ the excursion away from $(0,0)$ of $((\rho_t,\mathcal{N}_t) : t\geq 0)$ in the interval (α_j,β_j):

$$\begin{cases}\rho^j_s = \rho_{\alpha_j+s} & 0 < s < \beta_j-\alpha_j \\ \rho^j_s = 0 & s \geq \beta_j - \alpha_j\end{cases}, \qquad \begin{cases}\mathcal{N}^j_s = \mathcal{N}_{\alpha_j+s} & 0 < s < \beta_j-\alpha_j \\ \mathcal{N}^j_s = 0 & s \geq \beta_j - \alpha_j\end{cases}.$$

Then, the Markov process $((\rho_t,\mathcal{N}_t) : t\geq 0)$ can be rewritten as

$$(\rho_t,\mathcal{N}_t) = \sum_{j\in J}\mathbf{1}_{\{\alpha^j<t<\beta^j\}}(\rho^j_{t-\alpha^j},\mathcal{N}^j_{t-\alpha^j}), \quad t\geq 0. \tag{6.1}$$

The point process in $\mathbb{R}_+\times\mathbb{D}(\mathbb{R}_+,\mathscr{M}_f(\mathbb{R}_+)\times\mathscr{M}(\mathbb{R}^2_+))$ given by

$$\mathbf{M}(\mathrm{d}\ell,\mathrm{d}\rho,\mathrm{d}\mathcal{N}) := \sum_{j\in J}\delta_{(\ell^j,\rho^j,\mathcal{N}^j)}(\mathrm{d}\ell,\mathrm{d}\rho,\mathrm{d}\mathcal{N}), \tag{6.2}$$

where $(\ell^j = L^0_{\alpha_j})_{j\in J}$, is Poisson of intensity $\mathrm{d}x \otimes \mathbb{N}(\mathrm{d}\rho\,,\mathrm{d}\mathcal{N})$ with $\mathbb{N}(\mathrm{d}\rho\,,\mathrm{d}\mathcal{N}) = \mathbf{N}(\mathrm{d}\rho)Q^{H(\rho)}(\mathrm{d}\mathcal{N})$, $\mathbf{N}(\mathrm{d}\rho)$ the excursion measure of the exploration process and $Q^{H(\rho)}$ the conditional (probability) law of $(\mathcal{N}_t : t \geq 0)$ given ρ. These facts follow from standard excursion theory, or are established in Section 4.1.4 in [11] (in particular in what concerns the description of $\mathbb{N}$). [4]

Reciprocally, given a Poisson point process $\mathbf{M}$ of intensity $\mathrm{d}x \otimes \mathbb{N}(\mathrm{d}\rho\,,\mathrm{d}\mathcal{N})$ and atoms $(\ell^j, \rho^j, \mathcal{N}^j)_{j\in J}$, a snake process $((\rho_t, \mathcal{N}_t) : t \geq 0)$ is uniquely determined by the relation (6.1), with (α_j, β_j) defined in terms of $\mathbf{M}$ by

$$\beta_j := \sum_{k\in J:\ell^k \leq \ell^j} \zeta_k \quad \text{and } \alpha_j := \sum_{k\in J:\ell^k < \ell^j} \zeta_k\,,$$

with $\zeta_j := \inf\{s \geq 0 : \rho^j_s = 0\}$ the length of excursion j for each $j \in J$ (the fact that the measure $\mathrm{d}L^0_t = \mathbf{1}_{\{\rho_t = 0\}}\mathrm{d}L^0_t$ is singular with respect to $\mathrm{d}t$ is used to check this).

For a fixed height $a \geq 0$ we will also consider the process $(\rho^a_t, \mathcal{N}^a_t)_{t\geq 0}$ describing "what happens above a", that is, ρ^a_t is defined as in (1.9) and

$$\mathcal{N}^a_t(\mathrm{d}r, \mathrm{d}\nu) := \mathcal{N}_{\tau^a_t}(a + \mathrm{d}r, \mathrm{d}\nu),$$

with τ^a_t the stopping time defined in (1.8). We moreover denote by $(\rho^{(i)})_{i\in I}$ the excursions of the process ρ above height a and by $(\alpha^{(i)}, \beta^{(i)})_{i\in I}$ the corresponding excursion intervals. More precisely, for each $i \in I$, we set

$$\begin{cases} \langle \rho^{(i)}_s, f\rangle = \displaystyle\int_{(a,\infty)} \rho_{\alpha^{(i)}+s}(\mathrm{d}r) f(r-a) & 0 < s < \beta^{(i)} - \alpha^{(i)} \\ \qquad \rho^{(i)}_s = 0 & s \geq \beta^{(i)} - \alpha^{(i)}. \end{cases}$$

These excursions are in one-to-one correspondence to the excursions away from 0 of the process ρ^a occurring at cumulated local times $L^a_{\alpha^{(i)}} = L^a_{\beta^{(i)}}$ at level a. We also introduce the excursions of $\mathcal{N}$ above level a, "relative to a", given by [5]

$$\begin{cases} \mathcal{N}^{(i)}_s(\mathrm{d}r, \mathrm{d}\nu) = \mathcal{N}_{\alpha^{(i)}+s}(a + \mathrm{d}r, \mathrm{d}\nu) & 0 < s < \beta^{(i)} - \alpha^{(i)} \\ \qquad \mathcal{N}^{(i)}_s = 0 & s \geq \beta^{(i)} - \alpha^{(i)} \end{cases}.$$

Remark 6.1. Each excursion $\mathcal{N}^{(i)}$ away from 0 corresponds to a segment of a unique "parent excursion" $\mathcal{N}^j$ such that $(\alpha^{(i)}, \beta^{(i)}) \subset (\alpha^j, \beta^j)$.

Thus, $(\rho^{(i)}, \mathcal{N}^{(i)})_{i\in I}$ are exactly the excursions of the process $((\rho^a_t, \mathcal{N}^a_t) : t \geq 0)$ away from $(0, 0)$. The next result follows from [12, Prop 3.1] and the snake property (but see Appendix A.3 for the sketch of a direct proof, adapted from the one of Proposition 4.2.3 in [11]) :

Lemma 6.1 (Snake-excursion process above a given level). *For each $a \geq 0$, conditionally on the sigma field $\mathcal{E}_a$ defined in* (4.11), *the point process in $\mathbb{R}_+ \times \mathbb{D}(\mathbb{R}_+, \mathcal{V})$ given by*

$$\mathbf{M}^a(\mathrm{d}\ell, \mathrm{d}\rho\,, \mathrm{d}\mathcal{N}) := \sum_{i\in I} \delta_{(\ell^{(i)}, \rho^{(i)}, \mathcal{N}^{(i)})}(\mathrm{d}\ell, \mathrm{d}\rho\,, \mathrm{d}\mathcal{N}), \tag{6.3}$$

where $\ell^{(i)} = L^a_{\alpha^{(i)}}$ for all $i \in I$, has the same law as the process (6.2) *and is thus independent of $\mathcal{E}_a$. Consequently, $((\rho^a_t, \mathcal{N}^a_t) : t \geq 0)$ has the same law as $((\rho_t, \mathcal{N}_t) : t \geq 0)$ and is independent of $\mathcal{E}_a$.*

---

[4] Notice that results in [11] are stated in terms of the process $((\rho_t, W_t) : t \geq 0)$ with W_t corresponding here to the increasing $\mathcal{M}_{at}(\mathbb{R}_+)$-valued path $W_t = (h \mapsto \mathcal{N}_t([0, h], d\nu) : h \in [0, H_t)\,)$, which was denoted $\xi^{(t)}$ in Section 3.

[5] In the notation of [11], $\mathcal{N}^{(i)}$ would correspond to the "increment" of an excursion of the path-valued process W_t.

6.2 Proofs of Proposition 4.2 and Lemma 4.2

The following result extending Lemma 6.1 is a cornerstone of this paper. Consider an intensity process ϑ and for a given height $a > 0$ define:

$$\vartheta^a(t,h) := \mathbf{1}_{[0,a]}(h)\vartheta(t,h) \text{ for all } (t,h) \in \mathbb{R}_+^2.$$

Denote also

$$I^{\vartheta^a} := \{i \in I \,:\, m^{\vartheta^a}_{\alpha^{(i)}} = 0\} = \{i \in I \,:\, m^{\vartheta^a}_{\alpha^{(i)}}([0,a)) = 0\}.$$

By the semi-snake property if m^{ϑ^a}, for every $i \in I$ one has $m^{\vartheta^a}_s = 0 \forall s \in [\alpha^{(i)}, \beta^{(i)}] = 0$, thus I is the set of excursions above $a > 0$ whose parent excursion $\mathcal{N}^j$ has no marks below that level in the time interval $(\alpha^{(i)} - \alpha^j, \beta^{(i)} - \alpha^j)$. We have

Lemma 6.2 (Pruning below a fixed level). *The process* $(L^a_{T^a_x}(m^{\vartheta^a}) : x \geq 0)$ *is measurable with respect to the sigma-field* $\mathcal{B}(\mathbb{R}_+) \otimes \mathcal{E}_a$*, where* $\mathcal{E}_a$ *was defined in* (4.11)*. Moreover, conditionally on* $\mathcal{E}_a$*, the point process in* $\mathbb{R}_+ \times \mathbb{D}(\mathbb{R}_+, \mathcal{V})$ *given by*

$$\sum_{i \in I^{\vartheta^a}} \delta_{(L^a_{\alpha^{(i)}}(m^{\vartheta^a}), \rho^{(i)}, \mathcal{N}^{(i)})}(\mathrm{d}\ell, \mathrm{d}\rho\,, \mathrm{d}\mathcal{N}), \tag{6.4}$$

with $L^a_t(m^{\vartheta^a})$ *the* m^{ϑ^a}*-pruned local time at height* $a > 0$ *and time* $t \geq 0$*, is equal in law to the process* (6.2) *stopped at* $\ell = L^a_\infty(m^{\vartheta^a})$*.*

Remark 6.2. It is not hard to see that the marked exploration process associated with the excursion process (6.4) is given by

$$\left(\widehat{\rho}_t, \widehat{\mathcal{N}}_t \,: t \in [0, A^{\vartheta^a}_\infty)\right) = \left(\rho_{\widehat{C}_t}, \mathcal{N}_{\widehat{C}_t} \,: t \in [0, A^{\vartheta^a}_\infty)\right)\,,$$

where $\widehat{C}_t := \inf\left\{s > 0 \,:\, A^{\vartheta^a}_s > t\right\}$ and $A^{\vartheta^a}_s := \int_0^s \mathbf{1}_{\{m^{\vartheta^a}_r([0,a))=0\}} \mathrm{d}A^a_r$. Indeed, by the semi-snake property, $\mathbf{1}_{\{m^{\vartheta}_u([0,a))=0\}}$ is constant on the increase intervals of $A^a_u = \int_0^u \mathbf{1}_{\{H_r > a\}}\mathrm{d}r$, hence the only excursions above a contributing to the integral in the definition of $\widehat{C}_t$ are those in the set I^{ϑ^a}.

Lemma 6.2 will follow from Lemma 6.1 and an elementary fact about Poisson processes in $\mathbb{R}_+$, proved in Appendix A.4 for completeness:

Lemma 6.3. *Let* $(N_x : x \geq 0)$ *be a standard Poisson process of parameter* $\lambda > 0$ *with respect to a given filtration* $(\mathcal{K}_\ell)_{\ell \geq 0}$ *and* $E \subset \mathbb{R}_+$ *be a random set such that* $\mathbf{1}_E$ *is* $\mathcal{B}(\mathbb{R}_+) \otimes \mathcal{K}_0$*-measurable. For each* $x, y \geq 0$ *set* $\varphi_x := \int_0^x \mathbf{1}_E(\ell)\mathrm{d}\ell$ *and define the stopping time* $\phi_y := \inf\{x \geq 0 : \varphi_x > y\} \leq \infty$*. Then, conditionally on* $\mathcal{K}_0$ *,* $N^E := \left(\int_0^{\phi_x} \mathbf{1}_E(\ell)N(\mathrm{d}\ell) \,:\, x \geq 0\right)$ *is a standard* $(\mathcal{K}_{\phi_x})_{x \geq 0}$*-Poisson process of parameter* λ *stopped at* $\int_0^\infty \mathbf{1}_E(\ell)\mathrm{d}\ell$*.*

Proof of Lemma 6.2. In the proof we write $\vartheta^a = \vartheta$ for simplicity. As in the proof of Proposition 4.2.3 in [11] we introduce $\tilde{L}^a_t := L^a_{\tilde{\tau}^a_t}$ with $\tilde{\tau}^a_t$ defined above after (4.11), and its right-continuous inverse $\gamma^a(r) := \inf\{s \geq 0 \,:\, \tilde{L}^a_s > r\}$. Let us rewrite the m^ϑ-pruned local time at level a in terms of the corresponding local time units. Using the semi-snake property of m^ϑ in the second equality, we have

$$\begin{aligned} L^a_{T^a_x}(m^\vartheta) &= \int_0^{T^a_x} \mathbf{1}_{\{m^\vartheta_v = 0\}} \mathrm{d}L^a_v = \int_0^{\tilde{A}^a_{T^a_x}} \mathbf{1}_{\{m^\vartheta_{\tilde{\tau}^a_u} = 0\}} \mathrm{d}\tilde{L}^a_u \\ &= \int_0^{\tilde{L}^a_{\tilde{A}^a_{T^a_x}}} \mathbf{1}_{\{m^\vartheta_{\tilde{\tau}^a_{\gamma^a(\ell)}}([0,a))=0\}} \mathrm{d}\ell = \int_0^x \mathbf{1}_{\{\mathbf{m}^\vartheta_\ell = 0\}} \mathrm{d}\ell \end{aligned}$$

where for all $\ell \geq 0$ we have set $\mathbf{m}_\ell^\vartheta := m^\vartheta_{\tilde{\tau}^a_{\gamma^a(\ell)}}([0,a))$. We have also used the changes of variables $v = \tilde{\tau}^a_u$ and $\ell = \tilde{L}^a_u$. The last equality above stems from the fact that, by definition of $\tilde{\tau}^a_s$ (as the right inverse of $\tilde{A}^a_t = \int_0^t \mathbf{1}_{\{H_s \leq a\}} \mathrm{d}s$) and since L^a_s does not vary on intervals where $H_s > a$, one has $\tilde{L}^a_{\tilde{A}^a_{T^a_x}} = L^a_{\tilde{\tau}^a_{\tilde{A}^a_{T^a_x}}} = L^a_{T^a_x}$. Given that $\tilde{\tau}^a_t < \infty$ a.s. for each $t \geq 0$, (3.5) implies that

$$L^a_{\tilde{\tau}^a_s} = \lim_{\epsilon \to 0} \epsilon^{-1} \int_0^s \mathbf{1}_{\{a-\epsilon < H_{\tilde{\tau}^a_r}\}} \mathrm{d}r \quad \text{in probability uniformly on } s \text{ in compact sets.}$$

It follows that the process $(\mathbf{1}_{\{\mathbf{m}^\vartheta_\ell = 0\}} : \ell \geq 0)$ is $\mathcal{B}(\mathbb{R}_+) \otimes \mathcal{E}_a$-measurable. Thus, $(L^a_{T^a_x}(m^\vartheta) : x \geq 0)$ is $\mathcal{B}(\mathbb{R}_+) \otimes \mathcal{E}_a$-measurable as claimed.
We now introduce the filtration $\sigma\left(\{\mathbf{M}^a([0,x], \mathrm{d}\rho, \mathrm{d}\mathcal{N}) : 0 \leq x \leq \ell\} \vee \mathcal{E}_a\right)_{\ell \geq 0}$ (with $\mathbf{M}^a$ given in (6.3)) and its right-continuous completion denoted $(\mathcal{K}^a_\ell)_{\ell \geq 0}$. For each Borel set $S \subset \mathbb{D}(\mathbb{R}_+, \mathcal{V})$ with $\mathbb{N}(S) < \infty$, define a $(\mathcal{K}^a_\ell)_{\ell \geq 0}$- Poisson process in $\mathbb{R}_+$ by

$$N^{a,S}([0,\ell]) := \mathbf{M}^a([0,\ell] \times S), \quad \ell \geq 0.$$

Setting $\phi_y = \inf\{x > 0 : \int_0^x \mathbf{1}_{E^\vartheta(\ell)} d\ell > y\}$ with $E^\vartheta := \{\ell \in \mathbb{R}_+ : \mathbf{m}^\vartheta_\ell = 0\}$, Lemma 6.3 yields that conditionally on $\mathcal{K}^a_0$,

$$N^{E^\vartheta}(S) := \left(\int_0^{\phi_x} \mathbf{1}_{E^\vartheta}(\ell) N^{a,S}(\mathrm{d}\ell) : x \geq 0 \right)$$

is a Poisson process in $\mathbb{R}_+$ of parameter $\mathbb{N}(S)$, with respect to the time changed filtration $(\mathcal{K}^a_{\phi_l \geq 0})_{\ell \geq 0}$. Moreover, the processes $N^{E^\vartheta}(S_1), .., N^{E^\vartheta}(S_n)$ do not have simultaneous jumps if the sets $S_1, ..., S_n$ are disjoint, hence they are independent from each other conditionally on $\mathcal{K}^a_0$. We conclude that, conditionally on $\mathcal{K}^a_0$, the point process $\mathbf{M}^{a,\vartheta}$ on $\mathbb{R}_+ \times \mathbb{D}(\mathbb{R}_+, \mathcal{V})$ defined by

$$\mathbf{M}^{a,\vartheta}([0,x] \times S) = \int_{[0,\phi_x] \times S} \mathbf{1}_{E^\vartheta}(\ell) \mathbf{M}^a(\mathrm{d}\ell, \mathrm{d}\rho, \mathrm{d}\mathcal{N})$$

is Poisson with intensity $\mathrm{d}x \otimes \mathbb{N}(\mathrm{d}\rho, \mathrm{d}\mathcal{N})$. It is not hard to see that this is exactly the process (6.4). $\square$

We are ready to give the

Proof of Proposition 4.2. The fact that for each $(k,n) \in \mathbb{N}^2$ the process (4.12) is $\mathcal{F}_{\overline{T}_{(n+1)\delta}}$-measurable is immediate. For all the remaining properties, we proceed by induction on $k \in \mathbb{N}$.
In the case $k = 0$, the first assertion is obvious since for every $n \in \mathbb{N}$ one has $L^0_{\overline{T}^0_{(n+1)\delta}}(\varepsilon,\delta) - L^0_{\overline{T}^0_{n\delta}}(\varepsilon,\delta) = L^0_{T^0_{(n+1)\delta}}(\varepsilon,\delta) - L^0_{T^0_{n\delta}}(\varepsilon,\delta) = \delta$ a.s. and $\mathcal{E}_0$ is trivial. For the second assertion, we observe first that in a similar way as in the proof of Lemma 6.2, using (3.5) one shows that $L^h_{\tilde{\tau}^\varepsilon_s} = \lim_{\eta \to 0} \eta^{-1} \int_0^s \mathbf{1}_{\{\varepsilon - \eta < H_{\tilde{\tau}^\varepsilon_r} \leq h\}} \mathrm{d}r$ in probability uniformly on s in compact sets, for each $h \in [0, \varepsilon]$. Furthermore, in a similar way as in that proof we also get that

$$L^h_{T_x}(\varepsilon,\delta) = \int_0^{\check{L}^h_{\tilde{A}^\varepsilon_{T_x}}} \mathbf{1}_{\{m^{\varepsilon,\delta}_{\tilde{\tau}^\varepsilon_{\check{\gamma}(\ell)}}([0,h)) = 0\}} \mathrm{d}\ell,$$

where $s \mapsto \check{L}^h_s := L^h_{\tilde{\tau}^\varepsilon_s}$ and its right-continuous inverse denoted $\check{\gamma}(\ell)$ are both $\mathcal{B}(\mathbb{R}_+) \otimes \mathcal{E}_\varepsilon$-measurable. Since $\tilde{\tau}^\varepsilon_{\tilde{A}^\varepsilon_u} \geq u$ with equality if $u \geq 0$ is a strict right-increase point of $\tilde{A}^\varepsilon_u$, we deduce that

$$L^0_{\tilde{\tau}^\varepsilon_{\tilde{A}^\varepsilon_{T_x}}} = L^0_{T_x} = x \quad \text{and that} \quad \tilde{A}^\varepsilon_{T_x} = \inf\{s \geq 0 : L^0_{\tilde{\tau}^\varepsilon_s} > x\}.$$

The latter random variable is thus $\mathcal{E}_\varepsilon$-measurable. It follows that $L^h_{T^0_x}(\varepsilon,\delta)$ is $\mathcal{E}_\varepsilon$-measurable for every $x \geq 0$ and $h \in [0,\varepsilon]$ as required. The fact that $\left(L^h_{T^0_{n\delta+z}}(\varepsilon,\delta) - L^h_{T^0_{n\delta}}(\varepsilon,\delta) : h \in [0,\varepsilon], z \in [0,\delta]\right)$ is $\mathcal{F}_{T^0_{(n+1)\delta}}$-measurable is obvious, and it has the asserted conditional law thanks Proposition 1.2 and the strong Markov property of $(\rho,\mathcal{N})$ applied at $\mathcal{F}_{T^0_{n\delta}}$. This completes the case $k=0$.
For the inductive step, assume the statements are valid for all integer smaller than or equal to $(k-1)$. Since for each $n \in \mathbb{N}$,

$$L^{k\varepsilon}_{\overline{T}^{k\varepsilon}_{(n+1)\delta}}(\varepsilon,\delta) - L^{k\varepsilon}_{\overline{T}^{k\varepsilon}_{n\delta}}(\varepsilon,\delta) = \delta \wedge (L^{k\varepsilon}_\infty(\varepsilon,\delta) - n\delta)_+$$

and $L^{k\varepsilon}_\infty = \lim\limits_{l\to\infty} L^{k\varepsilon}_{\overline{T}^{(k-1)\varepsilon}_{l\delta}}$ is $\mathcal{E}_{k\varepsilon}$-measurable by the induction hypothesis, the first property for the integer k is immediate. We consider next the marked exploration process $((\rho_t, m^{\vartheta^{k\varepsilon}}_t) : t \geq 0)$ with adapted intensity $\vartheta^{k\varepsilon}(t,h) := \vartheta^{\varepsilon,\delta}(t,h)\mathbf{1}_{\{h\leq k\varepsilon\}}$, that is, $\vartheta^{k\varepsilon} = (\vartheta^{\varepsilon,\delta})^{k\varepsilon}$ in the notation of the beginning of Section 6.2 with $a = k\varepsilon$. Also, denote respectively by $\widehat{L}^h_t$, $\widehat{T}_x$ and $\widehat{H}_t$ the local time processes at level $h \geq 0$, the inverse local time process at level 0 and the height process associated with the (possibly stopped) process $(\widehat{\rho},\widehat{\mathcal{N}}) = \left(\rho_{\widehat{C}_\cdot}, \mathcal{N}_{\widehat{C}_\cdot}\right)$ defined in Remark 6.2 (i.e. the marked exploration process pruned below level $a = k\varepsilon$). By already developed arguments using the approximation (3.5), we can check that for every $t,h \geq 0$, on the event that $\widehat{C}_t < \infty$ we have

$$\widehat{L}^h_t = \int_0^{\widehat{C}_t} \mathbf{1}_{\{m^{\varepsilon,\delta}_s([0,k\varepsilon))=0\}} \mathrm{d}L^{k\varepsilon+h}_s \tag{6.5}$$

a.s. (see the proof of Proposition 1.2 in A.1 for the details of a similar computation based on (3.4)). Thus, for each $y \in [0, L^{k\varepsilon}_\infty(\varepsilon,\delta))$ we have

$$L^{k\varepsilon+h}_{\overline{T}^{k\varepsilon}_y}(\varepsilon,\delta) = \int_0^{\overline{T}^{k\varepsilon}_y} \mathbf{1}_{\{m^{\varepsilon,\delta}_s([k\varepsilon,k\varepsilon+h))=0\}} \mathbf{1}_{\{m^{\varepsilon,\delta}_s([0,k\varepsilon))=0\}} \mathrm{d}L^{k\varepsilon+h}_s = \int_0^{A^{\vartheta^{k\varepsilon}}_{\overline{T}^{k\varepsilon}_y}} \mathbf{1}_{\{\widehat{m}^{\varepsilon,\delta}_u([0,h))=0\}} \mathrm{d}\widehat{L}^h_u$$

using the change of variable $s = \widehat{C}_u$, (6.5) and the notation $A^{\vartheta^{k\varepsilon}}_s = \int_0^s \mathbf{1}_{\{m^{\vartheta^{k\varepsilon}}_r([0,k\varepsilon))=0,\, H_r>k\varepsilon\}} \mathrm{d}r$ and

$$\widehat{m}^{\varepsilon,\delta}_u([0,h)) := m^{\varepsilon,\delta}_{\widehat{C}_u}([k\varepsilon, k\varepsilon+h))\,. \tag{6.6}$$

Notice now that since $H_{\overline{T}^{k\varepsilon}_y} = k\varepsilon$, with the notation used in Lemma 6.2 one has

$$A^{\vartheta^{k\varepsilon}}_{\overline{T}^{k\varepsilon}_y} = \sum_{i\in I^{\vartheta^{k\varepsilon}} : \, L^{k\varepsilon}_{\alpha^{(i)}}(m^{\vartheta^{k\varepsilon}}) \leq y} (\beta^{(i)} - \alpha^{(i)}) = \inf\{u \geq 0 : \widehat{L}^0_u > y\} = \widehat{T}_y. \tag{6.7}$$

Thus, we have

$$L^{k\varepsilon+h}_{\overline{T}^{k\varepsilon}_y}(\varepsilon,\delta) = \int_0^{\widehat{T}_y} \mathbf{1}_{\{\widehat{m}^{\varepsilon,\delta}_u([0,h))=0\}} \mathrm{d}\widehat{L}^h_u. \tag{6.8}$$

Therefore, thanks to Lemma 6.1 (with $a = k\varepsilon$), the construction of the processes (4.12) for the integer k can be done conditionally on $\mathcal{E}_{k\varepsilon}$ using the process $(\widehat{\rho},\widehat{\mathcal{N}})$ up to the time $\widehat{T}_y$ for every $y \leq L^{k\varepsilon}_\infty(\varepsilon,\delta)$, in the same way as in the case $k=0$ the processes (4.12) were constructed using the process $(\rho,\mathcal{N})$ up to each time T_y. In particular, the process (4.12) is measurable with respect to the sigma-field $\widehat{\mathcal{E}}_\varepsilon$, defined as in (4.11) in terms of the right-continuous inverse $\widehat{\tilde{\tau}}^\varepsilon_t$ of the process $\widehat{\tilde{A}}^\varepsilon_t := \int_0^t \mathbf{1}_{\{\widehat{H}_s\leq\varepsilon\}}\mathrm{d}s$. We can then check that $\widehat{\mathcal{E}}_\varepsilon \subseteq \mathcal{E}_{(k+1)\varepsilon}$, noting that $\widehat{C}_{\widehat{\tilde{\tau}}^\varepsilon_t}$ is the right-continuous inverse of the process

$\widetilde{\widehat{A}}^{\varepsilon}_{A_t^{\vartheta k\varepsilon}} = \int_0^t \mathbf{1}_{\{H_{\widehat{C}_{A_u^{\vartheta k\varepsilon}}} <(k+1)\varepsilon,\, m_u^{\vartheta}([0,k\varepsilon))=0,\, H_u>k\varepsilon\}}\mathrm{d}u = \int_0^t \mathbf{1}_{\{H_u<(k+1)\varepsilon,\, m_u^{\vartheta}([0,k\varepsilon))=0,\, H_u>k\varepsilon\}}\mathrm{d}u$, which yields $\widehat{C}_{\widetilde{\widehat{\tau}}^{\varepsilon}_t} = \tilde{\tau}^{(k+1)\varepsilon}_{\theta_t}$, with $\theta_t := \inf\left\{r>0 : \int_0^r \mathbf{1}_{\{m^{\vartheta}_{\tilde{\tau}_v^{(k+1)\varepsilon}}([0,k\varepsilon))=0,\, H_{\tilde{\tau}_v^{(k+1)\varepsilon}}>k\varepsilon\}}\mathrm{d}v > t\right\}$. Moreover, thanks to Lemma 6.1, the conditional law of the process (4.12) given $\mathcal{F}_{\overline{T}^{k\varepsilon}_{n\delta}} \bigvee \mathcal{E}_{k\varepsilon}$ is measurable with respect to $\widehat{\mathcal{F}}_{\widehat{T}_{(n+1)\delta}} \bigvee \mathcal{E}_{k\varepsilon}$, where $(\widehat{\mathcal{F}}_t)_{t\geq 0}$ is the filtration $(\mathcal{F}_{\widehat{C}_t})_{t\geq 0}$. Thus, the identification of that conditional law is done in a similar way as in the case $k=0$, reasoning in terms of $(\widehat{\rho}, \widehat{\mathcal{N}})$ conditionally on $\mathcal{E}_{k\varepsilon}$. This achieves the inductive step and concludes the proof.

□

We end this subsection with the

Proof of Lemma 4.2. To ease notation we write $m = m^{\vartheta}$. We furthermore define ϑ^{a+b} and $I^{\vartheta^{a+b}}$ as in the beginning of the present Section 6.2, but with $a+b$ instead of a.
Since $\breve{T}^{a+b}_{L_s^{a+b}(m)} \geq s$ for all $s \geq 0$, the result is obvious if $\breve{T}^a_x = \infty$. We thus assume that $\breve{T}^a_x < \infty$. Suppose moreover that $H_{t_0} = a+b+c$ for some $t_0 \in (\breve{T}^a_x, \breve{T}^{a+b}_{L^{a+b}_{\breve{T}^a_x}(m)})$ such that $m_{t_0}([0,a+b)) = 0$. Then, for some excursion $i \in I^{\vartheta^{a+b}}$ above level $a+b$ whose parent excursion above 0 has no marks below $a+b$, we must have $t_0 \in (\alpha^{(i)}, \beta^{(i)})$ and then $L^{a+b}_{\alpha^{(i)}}(m) = L^{a+b}_{t_0}(m)$. Since $L^{a+b}_{t_0}(m) \leq L^{a+b}_{\breve{T}^a_x}(m)$ by definition of $\breve{T}^{a+b}_{L^{a+b}_{\breve{T}^a_x}(m)}$, we get $\alpha^{(i)} \leq \breve{T}^a_x$, a contradiction since $\breve{T}^a_x < t_0$ and $H_{\breve{T}^a_x} = a < a+b$. Therefore, we get

$$\int_{\breve{T}^a_x}^{\breve{T}^{a+b}_{L^{a+b}_{\breve{T}^a_x}(m)}} \mathrm{d}L^{a+b+c}_t(m) = \int_{\breve{T}^a_x}^{\breve{T}^{a+b}_{L^{a+b}_{\breve{T}^a_x}(m)}} \mathbf{1}_{\{H_t=a+b+c\}}\mathrm{d}L^{a+b+c}_t(m) = 0$$

and we conclude the desired identity. □

6.3 Proof of Proposition 4.3

The remainder of this section is devoted to the proof of Proposition 4.3. The proof is quite technical and will need different types of estimates and localization with respect to different variables (which is one of the reasons why we only get convergence in probability). Next result provides a "continuity-type" estimate for pruned local times. Recall that the $(\mathcal{F}^{\rho}_t)$-stopping time T^M was defined in (3.8).

Lemma 6.4. *Let* $((\rho_t, m_t^{\vartheta_1}) : t \geq 0)$ *and* $((\rho_t, m_t^{\vartheta_2}) : t \geq 0)$*) be marked exploration process and* τ *be an arbitrary* $(\mathcal{F}^{\rho}_t)$*-stopping time.*

i) For $h \geq 0$ *we have that*

$$\mathbb{E}\left(\left|L^h_\tau(m^{\vartheta_1}) - L^h_\tau(m^{\vartheta_2})\right|\right) \leq \mathbb{E}\left(\int_0^\tau \mathrm{d}L^h_t \int_0^h |\vartheta_1(t,r) - \vartheta_2(t,r)|\,\mathrm{d}r\right).$$

ii) If moreover $\tau \leq T^M$ *a.s. for some* $M \geq 0$ *and for* $i = 1,2$ *there is* $\tilde{\vartheta}_i$ *such that* $\vartheta_i = g(L(m^{\tilde{\vartheta}_i}))$*, we have*

$$\mathbb{E}\left(\int_0^\tau \mathrm{d}L^h_t \int_0^h |\vartheta_1(t,r) - \vartheta_2(t,r)|\,\mathrm{d}r\right) \leq c(M)M\,\mathbb{E}\left(\int_0^\tau \mathbf{1}_{\{H_s\leq h\}} \int_0^{H_s} |\tilde{\vartheta}_1(s,r) - \tilde{\vartheta}_2(s,r)|\mathrm{d}r\mathrm{d}t\right)$$

where $c(M) > 0$ *is a Lipschitz constant of* g *in* $[0,M]$*.*

Proof. i) We have a.s. for all $t \geq 0$,

$$\left|L_t^h(m^{\vartheta_1}) - L_t^h(m^{\vartheta_2})\right| \leq \int_0^t \left|m_s^{\vartheta_1}([0,h)) - m_s^{\vartheta_2}([0,h))\right| \mathrm{d}L_s^h \leq \int_0^t f_h(s)\mathrm{d}L_s^h,$$

with $f_h(s) = \int_0^h \int_0^\infty \mathbf{1}_{\{\vartheta_1(s,r)\wedge\vartheta_2(s,r))<\nu\leq\vartheta_1(s,r)\vee\vartheta_2(s,r)\}}\mathcal{N}_s(\mathrm{d}r,\mathrm{d}\nu)$. Applying conditionally on $\mathcal{F}_s^\rho = \mathcal{G}_0^{(s)}$ the compensation formula we get for any $(\mathcal{F}_t^\rho)_{t\geq 0}$-stopping time τ that

$$\mathbb{E}\left(f_h(s)\mathbf{1}_{\{s\leq\tau\}}\middle|\mathcal{F}_s^\rho\right) \leq \int_0^h \mathbb{E}\left(|\vartheta_1(s,r) - \vartheta_2(s,r)|\middle|\mathcal{F}_s^\rho\right)\mathbf{1}_{\{s\leq\tau\}},$$

and the statement follows. For point ii), proceeding as in the first part of the proof of Lemma 3.4 we get

$$\mathbf{1}_{t\leq\tau}\int_0^h |\vartheta_1(t,r) - \vartheta_2(t,r)|\ \mathrm{d}r \leq c(M)\mathbf{1}_{t\leq\tau}\int_0^t \mathbf{1}_{\{H_s\leq h\}}\left|m_s^{\tilde{\vartheta}_1}([0,H_s)) - m_s^{\tilde{\vartheta}_2}([0,H_s))\right|\mathrm{d}s\,,$$

hence

$$\begin{aligned}\mathbb{E}\left(\int_0^\tau \mathrm{d}L_t^h \int_0^h |\vartheta_1(t,r) - \vartheta_2(t,r)|\ \mathrm{d}r\right) &\leq c(M)\mathbb{E}\left(\int_0^\tau \mathrm{d}L_t^h \int_0^t \mathbf{1}_{\{H_s\leq h\}}\left|m_s^{\tilde{\vartheta}_1}([0,H_s)) - m_s^{\tilde{\vartheta}_2}([0,H_s))\right|\right)\mathrm{d}s\\ &\leq c(M)M\int_0^\infty \mathbb{E}\left(\mathbf{1}_{\{s\leq\tau,\,H_s\leq h\}}\left|m_s^{\tilde{\vartheta}_1}([0,H_s)) - m_s^{\tilde{\vartheta}_2}([0,H_s))\right|\right)\mathrm{d}s.\end{aligned}$$

Conditioning on $\mathcal{F}_s^\rho$ inside the last expectation, and following the last lines in the proof of Lemma 3.4 yields the upper bound

$$c(M)M\int_0^\infty \mathbb{E}\left(\mathbf{1}_{\{s\leq\tau,\,H_s\leq h\}}\mathbb{E}\left(\int_0^{H_s}|\tilde{\vartheta}_1(s,r) - \tilde{\vartheta}_2(s,r)|\mathrm{d}r\middle|\mathcal{F}_s^\rho\right)\right)\mathrm{d}s$$

as well as the desired estimate. □

In the proof of Proposition 4.3 we will also need to compare accumulated pruned local times at heights that are not in the grid, with local times at heights in the grid right below. For local times pruned at fixed rate, this type of comparison can be deduced from a variant of the approximation (3.4). An extension to local-time dependent rates of the present framework is however not immediate, in part because the pruning rate is globally unbounded. We will thus need to localize such approximation argument with respect to accumulated local times, in order to deal with pruning rates taking values in compact intervals. In order to ensure that the localization parameter can be removed, while at the same time, making the grid parameters go to 0, we will moreover need quantitative information about the speed at which approximations of local times such as (3.4) converge. Such a result is the content of next lemma. Its technical proof, given in Appendix A.5, relies on a snake variant of L^2-Poisson calculus developed in [11] for the excursion of the exploration process. In a similar way as in that work, to avoid making additional integrability assumptions on the underlying Lévy process X_t, we need to also localize the exploration process with respect to its mass $\langle\rho_t, 1\rangle$. We thus introduce an additional parameter $K > 0$ and the stopping time

$$\tau^K := \inf\{s > 0\,:\,\langle\rho_s, 1\rangle \geq K\}.$$

Lemma 6.5 (Quantitative approximation of variably pruned local times at level 0**).** *Consider a càglàd function* $\Theta : \mathbb{R}_+ \to [0,\bar{\theta}]$ *with* $\bar{\theta} \geq 0$ *and the marked exploration process* $((\rho_t, m_t) : t \geq 0)$ *with* $m_t = m_t^\vartheta$ *associated with*

$$\vartheta(t,h) := \Theta(L_t^0) \quad \forall\,(t,h) \in \mathbb{R}_2^+\,.$$

a) There exists an explicit nonnegative function $(\varepsilon, \bar{\theta}) \mapsto \hat{\mathcal{C}}(\bar{\theta}, K, \varepsilon)$ going to 0 when $\varepsilon \to 0$ and increasing both in ε and $\bar{\theta}$, such that for all $x \geq 0$:

$$\mathbb{E}\left[\sup_{y\in[0,x]}\left|y - \frac{1}{\varepsilon}\int_0^{T_y} \mathbf{1}_{\{0<H_s\leq\varepsilon,\, m_s=0\}}\mathrm{d}s\right|\mathbf{1}_{\{\tau^K>T_x\}}\right] \leq \hat{\mathcal{C}}(\bar{\theta}, K, \varepsilon)(x+\sqrt{x}).$$

b) We deduce that for all $x \geq 0$,

$$\mathbb{E}\left[\sup_{t\in[0,T_x]}\left|L_t^{\varepsilon}(m) - L_t^0\right|\mathbf{1}_{\{\tau^K>T_x\}}\right] \leq \mathcal{C}(\bar{\theta}, K, \varepsilon)(x+\sqrt{x})$$

for some explicit nonnegative function $(\varepsilon, \bar{\theta}) \mapsto \mathcal{C}(\bar{\theta}, K, \varepsilon)$ with similar properties as $\hat{\mathcal{C}}(\bar{\theta}, K, \varepsilon)$.

Recall the notation $k_h = \sup\{k \in \mathbb{N} : k\varepsilon < h\}$ and $\overline{n}_t^k = \sup\{n \in \mathbb{N} : n\delta < L_t^{k\varepsilon}(\varepsilon, \delta)\}$ introduced for fixed $\varepsilon, \delta \geq 0$ in Sections 4.2 and 4.3. From the previous result we deduce

Lemma 6.6. *Let us fix real numbers $\varepsilon, \delta, K, M > 0$.*

a) For all $a > 0$,

$$\mathbb{E}\left[\sup_{t\leq T^M\wedge\tau^K}|L_t^{k_a\varepsilon}(\varepsilon,\delta) - L_t^a(\varepsilon,\delta)|\right] \leq \mathcal{C}(g(M), K, \varepsilon)(M+\sqrt{M}) + \Gamma(M,K)$$

where $\Gamma(M,K) = 2M\mathbb{P}(\tau^K \leq T_M)$.

b) For each $t \geq 0$,

$$\mathbb{E}\left[\mathbf{1}_{\{t<T^M\wedge\tau^K\}}\mathbf{1}_{\{H_t\leq a\}}\int_0^{H_t}|\vartheta^{\varepsilon,\delta}(t,h) - \vartheta^*(t,h)|\mathrm{d}h\right]$$
$$\leq c(M)a\left(\delta + \Gamma(M,K) + \mathcal{C}(g(M),K,\varepsilon)(M+\sqrt{M})\right)e^{c(M)t}.$$

Proof. a) We start noting that if $t \leq T^M$, we have $L_t^h(\varepsilon,\delta) \leq L_t^h \leq M$ for every $h \geq 0$. Thus, if $k_a = 0$, since $T^M \leq T_M$, we have

$$\sup_{t\leq T^M\wedge\tau^K}|L_t^{k_a\varepsilon}(\varepsilon,\delta) - L_t^a(\varepsilon,\delta)| \leq \sup_{t\leq T_M}|L_t^0 - L_t^a(\varepsilon,\delta)|\mathbf{1}_{\{\tau^K>T_M\}} + 2M\mathbf{1}_{\{\tau^K\leq T_M\}}$$

and the desired inequality follows from part b) of Lemma 6.5 with $\bar{\theta} = g(M)$. To prove the property for arbitrary $k_a = k$, we first observe that, since $T^M \leq T_M^{k\varepsilon}$,

$$\sup_{t\leq T^M\wedge\tau^K}|L_t^{k\varepsilon}(\varepsilon,\delta) - L_t^a(\varepsilon,\delta)| \leq \sup_{t\leq T_M^{k\varepsilon}}|L_t^{k\varepsilon}(\varepsilon,\delta) - L_t^a(\varepsilon,\delta)|\mathbf{1}_{\{\tau^K>T_M\}} + 2M\mathbf{1}_{\{\tau^K\leq T_M\}}, \tag{6.9}$$

so it enough to bound the first term on the right hand side of (6.9) by $\mathcal{C}(g(M),K,\varepsilon)(M+\sqrt{M})$ in order to obtain the desired inequality. To that end, consider again the processes $\widehat{L}_t^r$, $\widehat{T}_x$ and $\widehat{H}_t$ associated with $(\widehat{\rho}, \widehat{\mathcal{N}})$, the snake process pruned below level $k\varepsilon$ already used in the proof of Proposition 4.2. Observe that $\mathbf{1}_{\{\tau^K>T_M^{k\varepsilon}\}} \leq \mathbf{1}_{\{\widehat{\tau}^K>\widehat{T}_{L^{k\varepsilon}_{T_M^{k\varepsilon}}(\varepsilon,\delta)}\}}$, where $\widehat{\tau}^K := \inf\{s > 0 : \langle\widehat{\rho}_s, 1\rangle \geq K\}$, and that (by similar arguments as in the proof of Proposition 4.2) the supremum on the right-hand side of (6.9) satisfies

$$\sup_{t\leq T_M^{k\varepsilon}}|L_t^{k\varepsilon}(\varepsilon,\delta) - L_t^a(\varepsilon,\delta)| \leq \sup_{s\leq\widehat{T}_{L^{k\varepsilon}_{T_M^{k\varepsilon}}(\varepsilon,\delta)}}|\widehat{L}_s^0 - \int_0^s \mathbf{1}_{\{\widehat{m}_u^{\varepsilon,\delta}([0,\epsilon'))=0\}}d\widehat{L}_u^{\epsilon'}| = \sup_{s\leq\widehat{T}_{L^{k\varepsilon}_{T_M^{k\varepsilon}}(\varepsilon,\delta)}}|\widehat{L}_s^0 - \widehat{L}_s^{\epsilon'}(\widehat{m}^{\varepsilon,\delta})|,$$

where $\epsilon' = a - k\varepsilon \in [0,\varepsilon]$, and $\widehat{m}^{\varepsilon,\delta}$ is given in (6.6). Thanks to Lemma 6.2, when taking expectation to the first summand on the r.h.s. of (6.9), we can first condition on $\mathcal{E}_{k\varepsilon}$ and (conditionally) apply part b) of Lemma 6.5. The result then follows since $L^{k\varepsilon}_{T^{k\varepsilon}_M}(\varepsilon,\delta) \le M$ and $y \mapsto \mathcal{C}(g(y),K,\varepsilon)(y+\sqrt{y})$ is increasing.

b) Writing $\Delta^*_t := \int_0^{H_t} |\vartheta^{\varepsilon,\delta}(t,h) - \vartheta^*(t,h)|\mathrm{d}h$, we have

$$\mathbb{E}\left[\mathbf{1}_{\{t<T^M\wedge\tau^K\}}\mathbf{1}_{\{H_t\le a\}}\Delta^*_t\right] \le \mathbb{E}\left[\mathbf{1}_{\{t<T^M\wedge\tau^K\}}\mathbf{1}_{\{H_t\le a\}}\int_0^{H_t}|g(L^r_t(\varepsilon,\delta)) - g(L^r_t(m^*))|\,\mathrm{d}r\right] \tag{6.10}$$

$$+\,\mathbb{E}\left[\mathbf{1}_{\{t<T^M\wedge\tau^K\}}\mathbf{1}_{\{H_t\le a\}}\int_0^{H_t}\left|g\left(\overline{n}^{k_r}_t\delta\right) - g\left(L^{k_r\varepsilon}_t(\varepsilon,\delta)\right)\right|\mathrm{d}r\right] \tag{6.11}$$

$$+\,\mathbb{E}\left[\mathbf{1}_{\{t<T^M\wedge\tau^K\}}\mathbf{1}_{\{H_t\le a\}}\int_0^{H_t}\left|g\left(L^{k_r\varepsilon}_t(\varepsilon,\delta)\right) - g\left(L^r_t(\varepsilon,\delta)\right)\right|\mathrm{d}r\right]. \tag{6.12}$$

Taking into account the relations (3.12) and (4.9), the term on the r.h.s. of (6.10) is bounded by

$$c(M)\int_0^t \mathbb{E}\left[\mathbf{1}_{\{s<T^M\wedge\tau^K\}}\mathbf{1}_{\{H_s\le a\}}\Delta^*_s\right]\mathrm{d}s\,,$$

thanks to Lemma 3.4. From the definition of the integers $k_h = k_h(\varepsilon)$ and $\overline{n}^k_t = \overline{n}^k_t(\varepsilon,\delta)$, term (6.11) is bounded by $c(M)a\delta$. Finally, by part a) term (6.12) is bounded by

$$c(M)\mathbb{E}\left[\int_0^a \sup_{s\le T^M\wedge\tau^K}|L^{k_r\varepsilon}_s(\varepsilon,\delta) - L^r_s(\varepsilon,\delta)|\mathrm{d}r\right] \le c(M)\,a\left(\mathcal{C}(g(M),K,\varepsilon)(M+\sqrt{M}) + 2M\Gamma(M,K)\right).$$

The statement follows by Gronwall's lemma.

□

We can finally give the

Proof of Proposition 4.3. We fix $M > 0$ and $x \ge 0$ and consider the $(\mathcal{F}^\rho_t)-$stopping time

$$\tau = T \wedge \tau^K \wedge T_x$$

with $T, K \ge 0$ constants to be fixed later in terms of M. Thanks to the relations (3.12) and (4.9) we can apply inequality i) of Lemma 6.4 to get for all $a \ge 0$ that

$$\mathbb{E}\left[|L^a_{\tau\wedge T^M}(\varepsilon,\delta) - L^a_{\tau\wedge T^M}(m^*)|\right] \le \mathbb{E}\left[\int_0^{\tau\wedge T^M}\mathrm{d}L^a_s\int_0^a\left|g\left(\overline{n}^{k_r}_s\delta\right) - g\left(L^r_s(m^*)\right)\right|\mathrm{d}r\right],$$

from where

$$\mathbb{E}\left[|L^a_{\tau\wedge T^M}(\varepsilon,\delta) - L^a_{\tau\wedge T^M}(m^*)|\right] \le \mathbb{E}\left[\int_0^{\tau\wedge T^M}\mathrm{d}L^a_s\int_0^a\left|g\left(\overline{n}^{k_r}_s\delta\right) - g\left(L^{k_r\varepsilon}_s(\varepsilon,\delta)\right)\right|\mathrm{d}r\right] \tag{6.13}$$

$$+\,\mathbb{E}\left[\int_0^{\tau\wedge T^M}\mathrm{d}L^a_s\int_0^a\left|g\left(L^{k_r\varepsilon}_s(\varepsilon,\delta)\right) - g\left(L^r_s(\varepsilon,\delta)\right)\right|\mathrm{d}r\right] \tag{6.14}$$

$$+\,\mathbb{E}\left[\int_0^{\tau\wedge T^M}\mathrm{d}L^a_s\int_0^a|g(L^r_s(\varepsilon,\delta)) - g(L^r_s(m^*))|\mathrm{d}r\right]. \tag{6.15}$$

The right hand side of (6.13) is bounded by $c(M)M\,a\delta$, by construction of the process $L(\varepsilon,\delta)$ and definition of T^M. Term (6.14) is bounded by

$$\begin{aligned}
c(M)\,\mathbb{E}&\left[\int_0^{\tau\wedge T^M} \mathrm{d}L_s^a \int_0^a \sup_{u\le T^M\wedge\tau^K} |L_u^{k_r\varepsilon}(\varepsilon,\delta)-L_u^r(\varepsilon,\delta)|\mathrm{d}r\right]\\
&\le c(M)\,M\,\mathbb{E}\left[\int_0^a \sup_{u\le T^M\wedge\tau^K} |L_u^{k_r\varepsilon}(\varepsilon,\delta)-L_u^r(\varepsilon,\delta)|\mathrm{d}r\right]\\
&\le ac(M)M\left(\mathcal{C}(g(M),K,\varepsilon)(M+\sqrt{M})+\Gamma(M,K)\right)
\end{aligned}$$

where in the last inequality we used part a) of Lemma 6.6.
To bound (6.15), we use inequality ii) of Lemma 6.4 with the stopping time τ in that statement replaced by, in current proof's notation, $\tau\wedge T^M = T\wedge\tau^K\wedge T_x\wedge T^M \le T^M$, and with $\vartheta_1 = g(L(\varepsilon,\delta))$, $\vartheta_2 = g(L(m^*))$, $\tilde\vartheta_1 = \vartheta^{\varepsilon,\delta}$ and $\tilde\vartheta_2 = \vartheta^*$. This gives us the upper bound

$$c(M)M\,\mathbb{E}\left(\int_0^{T\wedge T_x} \mathbf{1}_{\{H_s\le a\}}\mathbf{1}_{\{s<T^M\wedge\tau^K\}}\int_0^{H_s}|\vartheta^{\varepsilon,\delta}(s,r)-\vartheta^*(s,r)|\mathrm{d}r\,\mathrm{d}t\right)$$

for expression (6.15). By Lemma 6.6 b), the latter is bounded by

$$a\,Tc(M)^2\,M\left(\delta+\Gamma(M,K)+\mathcal{C}(g(M),K,\varepsilon)(M+\sqrt{M})\right)e^{c(M)T}.$$

Bringing all together, we have shown that

$$\begin{aligned}
\mathbb{E}&\left[|L^a_{T_x\wedge(\tau^K\wedge T\wedge T^M)}(\varepsilon,\delta)-L^a_{T_x\wedge(\tau^K\wedge T\wedge T^M)}(m^*)|\right]\\
&\le ac(M)M(1+Tc(M))\left(\delta+\Gamma(M,K)+\mathcal{C}(g(M),K,\varepsilon)(M+\sqrt{M})\right)e^{c(M)T}.
\end{aligned}$$

We now choose for each $M>0$, $T:=M$. Since $\tau^K\to\infty$ when $K\to\infty$, for each $M>0$ we can moreover find some $K=K(M)$ going to ∞ with M and such that $\mathbb{P}(\tau^{K(M)}\le T_M)\le (c(M)M^3(1+M\,c(M)))e^{c(M)M})^{-1}$. With these choices, we have

$$c(M)M(1+M\,c(M))\Gamma(M,K(M))e^{c(M)M}\le M^{-1}\to 0$$

when $M\to\infty$, whereas the sequence of stopping times $\mathcal{T}_M := M\wedge\tau^{K(M)}\wedge T^M$ goes a.s. to ∞. Thus, for each $\eta>0$ and $M\ge 0$,

$$\begin{aligned}
\mathbb{P}\left[|L^a_{T_x}(\varepsilon,\delta)\right. & \left.-L^a_{T_x}(m^*)|>\eta\right]\\
&\le \mathbb{P}\left[|L^a_{T_x\wedge\mathcal{T}_M}(\varepsilon,\delta)-L^a_{T_x\wedge\mathcal{T}_M}(m^*)|>\eta\right]+\mathbb{P}(T_x>\mathcal{T}_M)\\
&\le \frac{ac(M)M(1+M\,c(M))}{\eta}\left(\delta+\Gamma(M,K(M))+\mathcal{C}(g(M),K(M),\varepsilon)(M+\sqrt{M})\right)e^{c(M)M}\\
&\quad +\mathbb{P}(T_x>\mathcal{T}_M).
\end{aligned}$$

Hence,

$$\begin{aligned}
\limsup_{(\varepsilon,\delta)\to(0,0)}\mathbb{P}\left[|L^a_{T_x}(\varepsilon,\delta)-L^a_{T_x}(m^*)|>\eta\right]\le&\frac{ac(M)M(1+M\,c(M))}{\eta}\Gamma(M,K(M))e^{c(M)M}\\
&+\mathbb{P}(T_x>\mathcal{T}_M).
\end{aligned}$$

Letting $M\to\infty$, we have shown that $\lim_{(\varepsilon,\delta)\to(0,0)}\mathbb{P}\left[|L^a_{T_x}(\varepsilon,\delta)-L^a_{T_x}(m^*)|>\eta\right]=0.$ □

Acknowledgments M.C.F.'s research was supported by a doctoral grant of BASAL-Conicyt Center for Mathematical Modeling (CMM) at University of Chile and a postdoctoral grant from the National Agency for Science and Technology Promotion, Argentina. She thanks also ICM Millennium Nucleus NC120062 (Chile) for travel support and a doctoral mobility grant of the French Embassy in Chile for a three months visit to Laboratoire de Probabilités et Modèles Aléatoires of University Paris 6 in 2012, as well as for hospitality during her stay. J.F. acknowledges partial support of ICM Millennium Nucleus NC120062 and BASAL-Conicyt CMM. The authors thank Jean-François Delmas for enlightening discussions at the beginning of this work and Victor Rivero and Juan Carlos Pardo for kindly providing some useful mathematical insights.

A Appendix

A.1 Proof of Proposition 1.2

Recall from [3, Theorem 0.3] that $(\rho_t^\theta : t \geq 0) := (\rho_{C_t^\theta} : t \geq 0)$, with C_t^θ the right-continuous inverse of $A_t^\theta := \int_0^t \mathbf{1}_{\{m_s^\theta = 0\}} \mathrm{d}s$, has the same law as the exploration process associated with a Lévy process of Laplace exponent (1.11). Denote by $\left(\bar{L}_s^a : t \geq 0\right)$ its local time at level a. Applying (3.4) to ρ^θ and performing the change of variable $C_r^\theta \mapsto u$, we deduce that

$$\bar{L}_t^a = \lim_{\epsilon \to 0} \epsilon^{-1} \int_0^{C_t^\theta} \mathbf{1}_{\{a < H_u \leq a+\epsilon,\, m_u^\theta = 0\}} \mathrm{d}u \quad \text{for all } t \geq 0,$$

in the $L^1(\mathbb{P})$ sense. Let us check that this limit is equal to $L^a_{C_t^\theta}(m^\theta)$. Since $C_t^\theta < \infty$ a.s. (see [3]), we deduce from (3.4) applied to the original exploration process $(\rho_t : t \geq 0)$ that, on the interval $[0, C_t^\theta]$, the finite measures $\epsilon^{-1} \mathbf{1}_{\{a-\epsilon < H_s \leq a\}} \mathrm{d}s$ converge weakly in probability towards $\mathrm{d}L_s^a$ as $\epsilon \to 0$. By Lemma 3.3 the function $s \mapsto \mathbf{1}_{\{m_s^\theta([0,H_s))=0\}}$ has a.s. at most countably many discontinuities, therefore a.s. it is continuous a.e. with respect to the measure $\mathbf{1}_{[0,C_t^\theta]}(s)\mathrm{d}L_s^a$. We thus deduce that:

$$L^a_{C_t^\theta}(m^\theta) = \mathbb{P} - \lim_{\epsilon \to 0} \epsilon^{-1} \int_0^{C_t^\theta} \mathbf{1}_{\{a < H_u \leq a+\epsilon,\, m_u^\theta = 0\}} \mathrm{d}u = \bar{L}_t^a. \tag{A.1}$$

By right-continuity in $t \geq 0$ of the first and third expressions, $\left(\bar{L}_t^a : t \geq 0\right)$ and $\left(L^a_{C_t^\theta}(m^\theta) : t \geq 0\right)$ are indistinguishable for each $a \geq 0$. In particular, if we set $\bar{T}_x = \inf\{s > 0 : \bar{L}_t^0 > x\}$, for each $a \geq 0$ we a.s. have

$$\bar{L}^a_{\bar{T}_x} = L^a_{C^\theta_{\bar{T}_x}}(m^\theta) = L^a_{T_x}(m^\theta)$$

since $\bar{L}^0_\cdot = L^0_{C^\theta_\cdot}(m^\theta)$ (by (A.1) with $a = 0$) and $L^0_\cdot(m^\theta) = L^0_\cdot$. The result follows from the above identities and Theorem 1.1 applied to the local times of ρ^θ.

A.2 Proof of estimate (5.7)

We follow ideas in the proofs of [10, Thm 2.1] and [13, Prop 3.1]. Define a function ϱ on $\mathbb{R}_+$ by $\varrho(x) := [\sigma + \int_0^1 r^2 \Pi(\mathrm{d}r)]\sqrt{x}$. Notice that for all $y \leq x$ one has

$$\int_0^\infty \mathrm{d}\nu \int_0^1 \Pi(\mathrm{d}r) \int_0^1 \frac{r^2 \mathbf{1}_{\{y<\nu<x\}}(1-t)}{\varrho(|x - y + tr\mathbf{1}_{\{y<\nu<x\}}|)^2} \mathrm{d}t \leq c := \frac{\int_0^1 r^2 \Pi(\mathrm{d}r)}{\left(\sigma + \int_0^1 r^2 \Pi(\mathrm{d}r)\right)^2}. \tag{A.2}$$

Consider a real sequence $\{a_j\}_{j\geq 1}$ defined by $a_0 = 1$ and $a_j = a_{j-1}\, e^{-j[\sigma + \int_0^1 r^2 \Pi(\mathrm{d}r)]^2}$, so that $a_j \searrow 0$ and $\int_{a_j}^{a_{j-1}} \varrho(z)^{-2}\mathrm{d}z = j$. For each $j \geq 1$, let ψ_j be a non-negative continuous function on $\mathbb{R}$ supported in (a_j, a_{j-1}) such that $0 \leq \psi_j(z) \leq 2j^{-1}\varrho(z)^{-2}$ and $\int_{a_j}^{a_{j-1}} \psi_j(z)\mathrm{d}z = 1$. Define also non-negative twice continuously differentiable functions ϕ_j on $\mathbb{R}$ by

$$\phi_j(x) = \int_0^{|x|} \mathrm{d}y \int_0^y \psi_j(z)\mathrm{d}z, \quad x \in \mathbb{R}.$$

Notice that for each $x \in \mathbb{R}$ it holds for all $j \geq 1$ that $0 \leq \phi_j'(x)\mathrm{sign}(x) \leq 1$, $\phi_j''(x) \geq 0$ and $\phi_j''(x)\sigma^2|x| \leq 2j^{-1}$. Moreover, we have $\phi_j(x) \nearrow |x|$ as $j \to \infty$.
For any $z, h, r, \nu, x, y \geq 0$ and a differentiable function f, set now $l(r, \nu;, x, y) := r\left[\mathbf{1}_{\{\nu < x\}} - \mathbf{1}_{\{\nu < y\}}\right]$, $\Delta_h f(z) := f(z+h) - f(z)$ and $D_h f(\zeta) := \Delta_h f(\zeta) - f'(\zeta)h$. Using (5.6) and Itô's formula, we get

$$\begin{aligned}
\phi_j(\zeta^{\varepsilon,\delta}_{t\wedge\tau_m}(x)) =& M_{t\wedge\tau_m} + \alpha_0 \int_0^{t\wedge\tau_m} \phi_j'(\zeta^{\varepsilon,\delta}_s(x))\zeta^{\varepsilon,\delta}_s(x)\mathrm{d}s \\
&+ \int_0^{t\wedge\tau_m} \int_0^\infty \int_1^\infty \Delta_{l(r,\nu;Z_s(x),Z^{\varepsilon,\delta}_s(x))}\phi_k(\zeta^{\varepsilon,\delta}_s(x))\mathrm{d}\nu\Pi(\mathrm{d}r)\mathrm{d}s \\
&+ \frac{\sigma^2}{2}\int_0^{t\wedge\tau_m} \phi_j''(\zeta^{\varepsilon,\delta}_s(x))\zeta^{\varepsilon,\delta}_s(x)\mathrm{d}s \\
&+ \int_0^{t\wedge\tau_m} \int_0^\infty \int_0^1 D_{l(r,\nu;Z_s(x),Z^{\varepsilon,\delta}_s(x))}\phi_k(\zeta^{\varepsilon,\delta}_s)\mathrm{d}\nu\Pi(\mathrm{d}r)\mathrm{d}s \\
&- \int_0^{t\wedge\tau_m} \phi_j'(\zeta^{\varepsilon,\delta}_s(x))\left[G(Z_s(x)) - G(Z^{\varepsilon,\delta}_s(x))\right]\mathrm{d}s \\
&- \int_0^{t\wedge\tau_m} \sum_{n=0}^{n_x^{k_s\varepsilon}} \int_0^{Z^{\varepsilon,\delta}_{k_s\varepsilon}(x)} \phi_j'(\zeta^{\varepsilon,\delta}_s(x))\mathbf{1}_{\{n\delta < u \leq (n+1)\delta\}}\left[g(u) - g(n\delta)\right]\mathrm{d}_u Z_{k_s\varepsilon,s}(u)\mathrm{d}s.
\end{aligned} \tag{A.3}$$

where $(M_{t\wedge\tau_m})_{t\geq 0}$ is a martingale. By properties of ϕ_j', the integrands in the first and second lines are respectively bounded by $|\alpha_0||Z_s(x) - Z^{\varepsilon,\delta}_s(x)|$ and $\int_1^\infty r\Pi(\mathrm{d}r)|Z_s(x) - Z^{\varepsilon,\delta}_s(x)|$. The fact that

$$0 \leq \int_0^\infty \int_0^1 D_{l(r,\nu;x,y)}\phi_j(x-y)\mathrm{d}\nu\Pi(\mathrm{d}r) \leq \frac{2c}{j}$$

following from estimate (A.2), together with the mentioned properties of ϕ_j'' ensure that the terms on the third and fourth lines vanish when $j \to \infty$. Taking expectation in (A.3) and letting $j \to \infty$, we get the desired bound, noting that $g(m)$ is a Lipschitz constant for the function G in $[0, m]$.

A.3 Proof of Lemma 6.1

In a similar way as for the process $((\rho_t, \mathcal{N}_t) : t \geq 0)$, the trajectories of the process $((\rho_t^a, \mathcal{N}_t^a) : t \geq 0)$ are determined in a unique (measurable) way from the atoms of the point process (6.3). It is therefore enough to establish the first claim. To do so, one easily adapts first the arguments of the proof of Proposition 4.2.3 in [11] in order to prove that, under the excursion measure $\mathbb{N}$, the process

$$\sum_{i\in I^j} \delta_{(\ell^{(i)}, \rho^{(i)}, \mathcal{N}^{(i)})}, \tag{A.4}$$

with $I_j := \{i \in I : (\alpha^{(i)}, \beta^{(i)}) \subset (\alpha^j, \beta^j)\}$ denoting the sub-excursions above level a of the excursion away from 0 labeled j, is conditionally on $\mathcal{E}_a$ a Poisson point process of intensity $\mathrm{d}x \mathbf{1}_{[L^a_{\alpha_j}, L^a_{\beta_j}]} \otimes \mathbb{N}(\mathrm{d}\rho\,, \mathrm{d}\mathcal{N})$ (our superscripts "(i)" correspond to superscripts "i" in the statement of [11]). The only difference is that, in the computation analogous to the one at end of that proof, we must consider here test functions depending also on the components $\ell^{(i)}$ of the atoms, and depending on the excursions of the spatial component above level a only through their increments respect to their values at that a. Since I is equal to the disjoint union $\bigcup_{j \in J} I_j$, one then concludes applying conditionally on $\mathcal{E}_a$ the additivity of Poisson point measures.

A.4 Proof of Lemma 6.3

By standard properties of Poisson processes, for any nonnegative predictable process f and stopping time τ in the given filtration, it holds that $\mathbb{E}\left[e^{-u\int_0^{\tau\wedge t} f(\ell)N(\mathrm{d}\ell) + \lambda\int_0^{\tau\wedge t}(1-e^{-uf(\ell)})\mathrm{d}\ell} | \mathcal{K}_0\right] = 1$ a.s. for all $u \geq 0$ and $t \geq 0$. If moreover τ is such that $\mathbb{E}\left[e^{\lambda\int_0^{\tau}(1-e^{-uf(\ell)})\mathrm{d}\ell}\right] < \infty$, by dominated convergence one gets after letting $t \to \infty$ that $\mathbb{E}\left[e^{-u\int_0^{\tau} f(\ell)N(\mathrm{d}\ell) + \lambda\int_0^{\tau}(1-e^{-uf(\ell)})\mathrm{d}\ell} | \mathcal{K}_0\right] = 1$ a.s. Now, from our assumptions and by a monotone class argument, we can check that $\mathbf{1}_E$ is a predictable process. Since $e^{\lambda\int_0^{\phi_x}(1-e^{-u\mathbf{1}_E(\ell)})\mathrm{d}\ell} = e^{\lambda\int_0^{\phi_x}(1-e^{-u})\mathbf{1}_E(\ell)\mathrm{d}\ell} = e^{\lambda(1-e^{-u})(x\wedge\int\mathbf{1}_E(\ell)\mathrm{d}\ell)} \leq e^{\lambda(1-e^{-u})x}$, from the previous we conclude that for all $x \geq 0$,

$$\mathbb{E}\left[e^{-u\int_0^{\phi_x}\mathbf{1}_E(\ell)N(\mathrm{d}\ell)} | \mathcal{K}_0\right] = e^{-\lambda x(1-e^{-u})} \text{ a.s.}$$

Applying conditionally on $\mathcal{K}_{\phi_y}$ this argument to increments $\int_{\phi_y}^{\phi_x}\mathbf{1}_E(\ell)N(\mathrm{d}\ell)$, the proof is complete.

A.5 Proof of Lemma 6.5

a) The proof is inspired by that of Lemma 1.3.2 in [11]. We have

$$\begin{aligned}
&\mathbb{E}\left(\sup_{y\in[0,x]}\left|y - \frac{1}{\varepsilon}\int_0^{T_y}\mathbf{1}_{\{0<H_s\leq\varepsilon,\, m_s=0\}}\mathrm{d}s\right|\mathbf{1}_{\{\tau^K > T_x\}}\right) \\
&\quad\leq \mathbb{E}\left(\sup_{y\in[0,x]}\left|\frac{1}{\varepsilon}\int_0^{T_y}\mathbf{1}_{\{0<H_s\leq\varepsilon, m_s=0, \langle\rho_s,1\rangle\leq K\}}\mathrm{d}s - \frac{1}{\varepsilon}\mathbb{E}\left(\int_0^{T_y}\mathbf{1}_{\{0<H_s\leq\varepsilon, m_s=0, \langle\rho_s,1\rangle\leq K\}}\mathrm{d}s\right)\right|\right) \\
&\qquad\qquad + \mathbb{E}\left(\sup_{y\in[0,x]}\left|\frac{1}{\varepsilon}\mathbb{E}\left(\int_0^{T_y}\mathbf{1}_{\{0<H_s\leq\varepsilon, m_s=0, \langle\rho_s,1\rangle\leq K\}}\mathrm{d}s\right) - y\right|\right)
\end{aligned} \tag{A.5}$$

The time integral in the above expressions can be written in terms of the excursion point process (6.2) as follows:

$$\int_0^{T_y}\mathbf{1}_{\{0<H_s\leq\varepsilon, m_s=0,\, \langle\rho_s,1\rangle\leq K\}}\mathrm{d}s = \sum_{j\in J_y}\int_0^{\zeta^j}\mathbf{1}_{\{0<H^j_s\leq\varepsilon, m^j_s=0,\, \langle\rho^j_s,1\rangle\leq K\}}\mathrm{d}s\,, \tag{A.6}$$

where $J_y := \{j \in J : \ell^j \leq y\}$, $H^j_s = H_{\alpha^j+s}$, $m^j_s = \mathcal{N}^j_s(\,\cdot \times [0, \Theta(\ell^j)))$ and ζ^j is the length of the excursion labelled j. By compensation, the desintegration $\mathbb{N}(\mathrm{d}\rho\,, \mathrm{d}\mathcal{N}) = \mathbf{N}(\mathrm{d}\rho)Q^{H(\rho)}(\mathrm{d}\mathcal{N})$ and the very definition of the snake $(\rho, \mathcal{N})$, we get

$$\begin{aligned}
&\mathbb{E}\left(\int_0^{T_y} \mathbf{1}_{\{0<H_s\le\varepsilon, m_s=0,\, \langle\rho_s,1\rangle\le K\}}\mathrm{d}s\right)\\
&=\int_0^y \mathrm{d}\ell\,\mathbb{N}\left(\int_0^\zeta \mathbf{1}_{\{0<H_s(\rho)\le\varepsilon, \mathcal{N}_s(\,\cdot\times[0,\Theta(\ell)))=0\,,\langle\rho_s,1\rangle\le K\}}\mathrm{d}s\right)\\
&=\int_0^y \mathrm{d}\ell\,\mathbf{N}\left(\int_0^\zeta e^{-\Theta(\ell)H_s(\rho)}\mathbf{1}_{\{0<H_s(\rho)\le\varepsilon\,,\langle\rho_s,1\rangle\le K\}}\mathrm{d}s\right),
\end{aligned}$$

with ζ the length of the canonical excursion and $(H_s(\rho) : 0 \le s \le \zeta)$ its height process. Thus, the second term in the r.h.s. of (A.5) is bounded by

$$\int_0^x \mathrm{d}\ell\,\mathbb{E}\left[\left|\varepsilon^{-1}\mathbf{N}\left(\int_0^\zeta e^{-\Theta(\ell)H_s(\rho)}\mathbf{1}_{\{0<H_s(\rho)\le\varepsilon\,,\langle\rho_s,1\rangle\le K\}}\mathrm{d}s\right)-1\right|\right].$$

Using Proposition 1.2.5 in [11] to compute the integral with respect to $\mathbf{N}$ for each $\ell \in [0, x]$, the latter expression is seen to be equal to

$$\int_0^x \mathrm{d}\ell\left[1-\frac{1}{\varepsilon}\int_0^\varepsilon e^{-\alpha b}e^{-\Theta(\ell)b}\mathbb{P}(S_b\le K)\mathrm{d}b\right]\le x\left[1-\frac{1-e^{-(\alpha+\bar\theta)\varepsilon}}{(\alpha+\bar\theta)\varepsilon}\mathbb{P}(S_\varepsilon\le K)\right] \tag{A.7}$$

where $(S_b)_{b\ge0}$ is a subordinator of Laplace exponent $\widehat{\psi}(\lambda) := \frac{\sigma^2}{2}\lambda + \int_0^\infty(1-e^{-\lambda})\Pi([r,\infty))\mathrm{d}r$ which does not depend on the drift coefficient α of the underlying Lévy process X. In particular, the expression on the r.h.s. of (A.7) goes to 0 with ε.

Let now $(\mathcal{Q}_\ell)_{\ell\ge0}$ denote the right continuous completion of the filtration $(\sigma(\mathbf{M}([0,x], d\rho, d\mathcal{N}) : 0 \le x \le \ell))_{\ell\ge0}$, with $\mathbf{M}$ the point process defined in (6.2). Since the first term on the r.h.s. of (A.5) is the expected supremum of the absolute value of a $(\mathcal{Q}_\ell)_{\ell\ge0}$-martingale, we can bound it using BDG inequality by some universal constant C_1 times

$$\sqrt{Var\left[\frac{1}{\varepsilon}\int_0^{T_x}\mathbf{1}_{\{0<H_s\le\varepsilon, m_s=0, \langle\rho_s,1\rangle\le K\}}\mathrm{d}s\right]}.$$

Written in terms of the excursion Poisson point process (6.2), the previous quantity reads

$$\sqrt{Var\left[\frac{1}{\varepsilon}\sum_{j\in J_x}\int_0^{\zeta_j}\mathbf{1}_{\{0<H^j_s\le\varepsilon, m^j_s=0,\, \langle\rho^j_s,1\rangle\le K\}}\mathrm{d}s\right]}$$

and can be estimated by the same arguments as in the proof of Lemma 1.3.2 of [11], as follows (see also the proof of Lemma 1.1.3 therein for details on related arguments):

$$\begin{aligned}
Var\left[\frac{1}{\varepsilon}\int_0^{T_x}\mathbf{1}_{\{0<H_s\le\varepsilon, m_s=0, \langle\rho_s,1\rangle\le K\}}\mathrm{d}s\right] &= \frac{x}{\varepsilon^2}\mathbb{N}\left(\left(\int_0^\zeta \mathbf{1}_{\{0<H_s\le\varepsilon, m_s=0, \langle\rho_s,1\rangle\le K\}}\mathrm{d}s\right)^2\right)\\
&\le \frac{x}{\varepsilon^2}\mathbf{N}\left(\left(\int_0^\zeta \mathbf{1}_{\{0<H_s\le\varepsilon, \langle\rho_s,1\rangle\le K\}}\mathrm{d}s\right)^2\right) \qquad \text{(A.8)}\\
&\le 2x\mathbb{E}(X_{L^{-1}(\varepsilon)}\wedge K),
\end{aligned}$$

where $\varepsilon \mapsto X_{L^{-1}(\varepsilon)}$ is a subordinator of Laplace exponent $\exp\left(-t\left(\widehat{\psi}(\lambda)+\alpha\right)\right)$. That is, the same subordinator S as above, but killed at an independent exponential time of parameter α. Thus, we have

$\mathbb{E}(X_{L^{-1}(\varepsilon)} \wedge K) \leq \mathbb{E}(S_\varepsilon \wedge K) + K(1 - e^{-\alpha\varepsilon}) \to 0$ as $\varepsilon \to 0$. The statement now follows by bringing together (A.5), (A.8) and (A.7) with the r.h.s. of the latter replaced by its supremum over $\varepsilon' \in [0, \varepsilon]$, which is an increasing function of ε going to 0 as $\varepsilon \to 0$, as required.

b) Observe first that by continuity of $s \mapsto L^0_s$

$$\forall t < T_x, \forall n \geq \frac{1}{x}, \exists y_n, z_n \in [\frac{1}{n}, x] \text{ such that } T_{y_n} \leq t \leq T_{z_n} \text{ and } z_n - \frac{1}{n} \leq L^0_t \leq y_n + \frac{1}{n}.$$

We deduce that

$$|L^0_t - L^\varepsilon_t(m)| \leq \begin{cases} y_n - L^\varepsilon_{T_{y_n}}(m) + \dfrac{1}{n} & \text{if } L^\varepsilon_t(m) < L^0_t \\ \\ L^\varepsilon_{T_{z_n}}(m) - z_n + \dfrac{1}{n} & \text{if } L^\varepsilon_t(m) > L^0_t \end{cases}.$$

Therefore, we have $\sup\limits_{t \in [0, T_x]} |L^0_t - L^\varepsilon_t(m)| \leq \sup\limits_{y \in [0,x]} \left| L^\varepsilon_{T_y}(m) - y \right|$ and it is enough to establish the required upper bound for the quantity

$$\mathbb{E}\left[\sup_{y \in [0,x]} \left| L^\varepsilon_{T_y}(m) - y \right| \mathbf{1}_{\{\tau^K > T_x\}} \right]. \tag{A.9}$$

We have

$$L^\varepsilon_{T_y}(m) - y = \left[L^\varepsilon_{T_y}(m) - \frac{1}{\varepsilon} \int_0^{T_y} \mathbf{1}_{\{\varepsilon < H_s \leq 2\varepsilon, m_s((0,\varepsilon)) = 0\}} \mathrm{d}s \right] + 2 \left[\frac{1}{2\varepsilon} \int_0^{T_y} \mathbf{1}_{\{0 < H_s \leq 2\varepsilon, m_s((0,\varepsilon)) = 0\}} \mathrm{d}s - y \right]$$
$$+ \left[y - \frac{1}{\varepsilon} \int_0^{T_y} \mathbf{1}_{\{0 < H_s \leq \varepsilon, m_s((0,\varepsilon)) = 0\}} \mathrm{d}s \right],$$

and the absolute value of the second term on the right hand side is bounded by

$$2 \left| \frac{1}{2\varepsilon} \int_0^{T_y} \mathbf{1}_{\{0 < H_s \leq 2\varepsilon, m_s((0,2\varepsilon)) = 0\}} \mathrm{d}s - y \right| + 2 \left| \frac{1}{2\varepsilon} \int_0^{T_y} \mathbf{1}_{\{0 < H_s \leq 2\varepsilon\}} \mathrm{d}s - y \right|,$$

thanks to the inequalities $\mathbf{1}_{\{0 < H_s \leq 2\varepsilon\}} \geq \mathbf{1}_{\{0 < H_s \leq 2\varepsilon, m_s((0,\varepsilon)) = 0\}} \geq \mathbf{1}_{\{0 < H_s \leq 2\varepsilon, m_s((0,2\varepsilon)) = 0\}}$. It follows from part a) that expression (A.9) is upper bounded by

$$\Big(2\hat{\mathcal{C}}(\bar{\theta}, K, 2\varepsilon) + 2\hat{\mathcal{C}}(0, K, \varepsilon) + \hat{\mathcal{C}}(\bar{\theta}, K, \varepsilon) \Big) (x + \sqrt{x})$$
$$+ \mathbb{E}\left[\sup_{y \in [0,x]} \left| L^\varepsilon_{T_y}(m) - \frac{1}{\varepsilon} \int_0^{T_y} \mathbf{1}_{\{\varepsilon < H_s \leq 2\varepsilon, m_s((0,\varepsilon)) = 0\}} \mathrm{d}s \right| \mathbf{1}_{\{\tau^K > T_x\}} \right] \tag{A.10}$$

and it only remains us to bound this last expectation. Notice to that end that the inner supremum can be written in terms of the Poisson excursions point process pruned below the level $a = \varepsilon$ as considered in (6.4), namely

$$\sum_{i \in I^{\vartheta^\varepsilon}} \delta_{(L^\varepsilon_{\alpha^{(i)}}(m), \rho^{(i)}, \mathcal{N}^{(i)})}, \tag{A.11}$$

where $I^{\vartheta^\varepsilon} = \{i \in I : m^{\vartheta^\varepsilon}_{\alpha^{(i)}}([0,\varepsilon)) = 0\}$. As before, let $\widehat{L}^0_t$, $\widehat{T}_x$, $\widehat{H}_t$ and $\widehat{\tau}^K$ denote the corresponding local time at 0, inverse local time at 0, height process, and the stopping time $\widehat{\tau}^K := \inf\{s > 0 : \langle \widehat{\rho}_s, 1 \rangle \geq K\}$, all of them now associated with $(\widehat{\rho}, \widehat{\mathcal{N}})$, the snake process associated with (A.11).

Then, writing in a similar way as in (A.6) the time integral in (A.10), that is, as a sum of integrals over (now) non marked excursion intervals above level ε, we get

$$L^{\varepsilon}_{T_y}(m) - \frac{1}{\varepsilon}\int_0^{T_y} \mathbf{1}_{\{\varepsilon < H_s \leq 2\varepsilon, m_s((0,\varepsilon))=0\}} \mathrm{d}s = \widehat{L}^{0}_{\widehat{T}_{L^{\varepsilon}_{T_y}(m)}} - \frac{1}{\varepsilon}\int_0^{\widehat{T}_{L^{\varepsilon}_{T_y}(m)}} \mathbf{1}_{\{0<\widehat{H}_s \leq \varepsilon\}} \mathrm{d}s.$$

Since $T_x \geq \widehat{T}_{L^{\varepsilon}_{T_x}(m)}$ and $\sup_{t \leq T_x} \langle \rho_t, 1\rangle \geq \sup_{t \leq \widehat{T}_{L^{a}_{T_x}}(m)} \langle \widehat{\rho}_t, 1\rangle$, the expectation in (A.10) is seen to be less than

$$\mathbb{E}\left[\sup_{z \in [0, L^{\varepsilon}_{T_x}(m)]} \left| z - \frac{1}{\varepsilon}\int_0^{\widehat{T}_z} \mathbf{1}_{\{0<\widehat{H}_s \leq \varepsilon\}} \mathrm{d}s \right| \mathbf{1}_{\{\widehat{\tau}^K > \widehat{T}_{L^{\varepsilon}_{T_x}(m)}\}}\right] \leq \hat{\mathcal{C}}(0, K, \varepsilon)\mathbb{E}\left[L^{\varepsilon}_{T_x}(m) + \sqrt{L^{\varepsilon}_{T_x}(m)}\right],$$

by applying, thanks to Lemma 6.2, the previous part a) (with $m = 0$ or equivalently $\bar{\theta} = 0$) conditionally on $\mathcal{E}_{\varepsilon}$. With the obvious bounds $L^{\varepsilon}_{T_x}(m) \leq L^{\varepsilon}_{T_x}$ a.s., $\mathbb{E}\left[\sqrt{L^{\varepsilon}_{T_x}}\right] \leq \sqrt{\mathbb{E}\left[L^{\varepsilon}_{T_x}\right]}$ and the equalities $\mathbb{E}(L^{\varepsilon}_{T_x}) = x\mathbf{N}\left(L^{\varepsilon}_{\zeta}\right) = xe^{-\alpha\varepsilon}$ following from Corollary 1.3.4 in [11], the desired result is seen to hold with $\mathcal{C}(\bar{\theta}, K, \varepsilon) = \left(2\hat{\mathcal{C}}(\bar{\theta}, K, 2\varepsilon) + 3\hat{\mathcal{C}}(0, K, \varepsilon) + \hat{\mathcal{C}}(\bar{\theta}, K, \varepsilon)\right)$.

A.6 Lamperti-type representation

Proposition A.1. *Let X_t be a Lévy process with regular Laplace exponent and assume that the locally bounded competition mechanisms g is such that $\lim_{z \to 0} \frac{g(z)}{z}$ exists. For each $x > 0$ there is a unique strong solution U_t to the SDE* (2.4)*. Moreover, if we set*

$$V_t := \begin{cases} U_{C_t} & \text{if } 0 \leq t < \eta_\infty, \\ 0 & \text{if } \eta_\infty < \infty \wedge t \geq \eta_\infty, \end{cases} \tag{A.12}$$

with C_t the right inverse of $\eta_t := \inf\left\{s > 0 : \int_0^s \frac{\mathrm{d}r}{U_r} > t\right\}$, there exists in some enlarged probability space a Brownian motion B^V and an independent Poisson point process N^V in $[0,\infty)^3$ with intensity measure $\mathrm{d}t \otimes \mathrm{d}\nu \otimes \Pi(\mathrm{d}r)$, such that

$$V_t = x - \alpha\int_0^t V_s \mathrm{d}s + \sigma\int_0^t \sqrt{V_s}\mathrm{d}B^V_s + \int_0^t\int_0^{V_{s-}}\int_0^\infty r\tilde{N}^V(\mathrm{d}s, \mathrm{d}\nu, \mathrm{d}r) - \int_0^t G(V_s)\mathrm{d}s, \tag{A.13}$$

for all $t \geq 0$. Last, pathwise uniqueness (and then also in law) holds for (A.13).

Since $(Z_t(x) : t \geq 0)$ in (1.5) satisfies (A.13) with the Brownian motion

$$B_t := \int_0^t \int_0^{Z_{s^-}(x)} \left(Z_{s^-}(x)\right)^{-\frac{1}{2}} W(\mathrm{d}s, \mathrm{d}u),$$

we conclude that $(Z_t(x) : t \geq 0)$ and $(V_t : t \geq 0)$ are equal in law.

Proof. Let B^X and N^X respectively denote a standard Brownian motion and a Poisson random measure on $[0,\infty)^2$ with intensity $\mathrm{d}s \otimes \Pi(\mathrm{d}r)$ such that $\mathrm{d}X_t = -\alpha\mathrm{d}t + \sigma\mathrm{d}B^X_t + \int_0^\infty r\widetilde{N}^X(\mathrm{d}t, \mathrm{d}r)$. Standard localization arguments using a sequence G^R of globally Lipschitz functions equal to G on $[0, R]$ (the local Lipschitz character of G following from the assumptions) show the existence of a unique strong solution U until some random explosion time. For each $R, K \geq 0$, set $\tau^R = \inf\{s \geq 0 : U^2_s \geq R^2\}$ and $\theta_K = \inf\{s \geq 0 : [U, U]_s \geq K\}$. Applying Itô' s formula to the solution of equation (2.4) with G_R

instead of G while keeping in mind the sign of G, we get using Gronwall's lemma that $\mathbb{E}(U^2_{t\wedge\theta_K\wedge\tau^R}) \leq c + c'K$ for some finite constants $c, c' > 0$ depending on t, x and the characteristics of X but not on G. Fatou's Lemma then yields $t \wedge \theta_K \leq \tau^\infty = \sup_{R\geq 0} \tau^R$ a.s., from where $\tau^\infty = \infty$ a.s. and

$$\mathrm{d}U_t = -\alpha \mathrm{d}t + \sigma \mathrm{d}B^X_t + \int_0^\infty r\widetilde{N}^X(\mathrm{d}t, \mathrm{d}r) - \frac{G(|U_t|)}{U_t}\mathrm{d}t. \tag{A.14}$$

Let us now set $T_0 := \inf\{t > 0 : U_t = 0\}$ and $T := \inf\{t > 0 : V_t = 0\} = \inf\{t > 0 : U_{C_t} = 0\} \wedge \eta_\infty$. As C_t is right-continuous, we have that $U_{C_T} = 0$, so that $C_T = T_0$ since $\eta_r = \int_0^{r\wedge T_0} \frac{\mathrm{d}s}{U_s} = \eta_\infty$ for all $r \geq T_0$. In order to show that the time-changed process $V = (U_{C_t} : t \geq 0)$ is solution of (A.13), we follow the arguments of Caballero *et al.* [9, Prop. 4] providing the existence of a standard Brownian motion B such that

$$\int_0^t \sqrt{V_s}\mathrm{d}B_s = B^X(C_t \wedge T_0) \tag{A.15}$$

and of a Poisson random measure N with intensity $\mathrm{d}s \otimes \mathrm{d}\nu \otimes \Pi(\mathrm{d}r)$ such that

$$\sum_{\{n:t^X_n<C_t\}} r^X_n = \sum_{\{n:t_n<t\}} \Delta_n = \int_0^t \int_0^{V_{s-}} \int_0^\infty r\mathbf{1}_{\{r\geq 1\}} N(\mathrm{d}s, \mathrm{d}\nu, \mathrm{d}r),$$

where $((\Delta_n, t_n) : n \in \mathbb{N})$ is a fixed but arbitrary labeling of the jump times and sizes of V and $((r^X_n, t^X_n) : n \in \mathbb{N})$ are the atoms of N^X. We have in an L^2 sense that

$$\lim_{\varepsilon\searrow 0}\left[\sum_{\{n:t_n<t\}} \Delta_n \mathbf{1}_{\{\Delta_n>\varepsilon\}} - \int_0^t V_s\mathrm{d}s \int_\varepsilon^\infty r\Pi(\mathrm{d}r)\right] = \lim_{\varepsilon\searrow 0}\left[\sum_{\{n:t^X_n<C_t\}} r^X_n \mathbf{1}_{\{r^X_n>\varepsilon\}} - \int_0^{C_t} \mathrm{d}s \int_\varepsilon^\infty r\Pi(\mathrm{d}r)\right], \tag{A.16}$$

so that the compensated measures satisfy

$$\int_0^t \int_0^{V_{s-}} \int_0^\infty r\widetilde{N}(\mathrm{d}s, \mathrm{d}\nu, \mathrm{d}r) = \int_0^{C_t} \int_0^\infty r\widetilde{N}^X(\mathrm{d}s, \mathrm{d}r). \tag{A.17}$$

Inserting the identities (A.15), (A.16) and (A.17) into equation (A.14), we deduce that

$$\mathrm{d}U_{C_t} = -\alpha\mathrm{d}C_t + \sigma\sqrt{V_t}\mathrm{d}B_t + \int_0^{V_{t-}} \int_0^\infty r\widetilde{N}(\mathrm{d}t, \mathrm{d}\nu, \mathrm{d}r) - cU_{C_t}\mathrm{d}C_t.$$

By (A.12), $\mathrm{d}C_t = V_t\mathrm{d}t$ and $\frac{G(|U_{C_t}|)}{U_{C_t}}\mathrm{d}C_t = G(|V_t|)\mathrm{d}t$, thus $V = (V_t : t \geq 0)$ is a solution of (A.13). Uniqueness for (A.13) follows from general results in [13]. $\square$

References

[1] R. ABRAHAM and J.-F. DELMAS. Fragmentation associated with Lévy processes using snake. *Probab. Theory Relat. Fields*, 141:113–154, 2008.

[2] R. ABRAHAM, J.-F. DELMAS, and H. HE. Pruning of CRT-sub-trees. *Stochastic Process. Appl.*, 125(4):1569–1604, 2015.

[3] R. ABRAHAM, J.-F. DELMAS, and G. VOISIN. Pruning a Lévy Continuum Random Tree. *Elec. J. of Probab.*, 15(46):1429–1473, 2010.

[4] R. ABRAHAM and L. SERLET. Poisson snake and fragmentation. *Elect. J. of Probab.*, 7:1–15, 2002.

[5] D. ALDOUS. The continuum random tree I. *Ann. Probab.*, 19:1–28, 1991.

[6] D. ALDOUS. The continuum random tree III. *Ann. Probab.*, 21:248–289, 1993.

[7] J. BERESTYCKI, L. DÖRING, L. MYTNIK, and L. ZAMBOTTI. On exceptional times for generalized Fleming-Viot processes with mutations. *Stoch. Partial Differ. Equ. Anal. Comput.*, 2(1):84–120, 2014.

[8] J. BERTOIN and J.-F. LE GALL. Stochastic flows associated to coalescent processes. *Probab. Relat. Fields*, 126:261–288, 2003.

[9] M. E. CABALLERO, A. LAMBERT, and G. URIBE BRAVO. Proof(s) of the Lamperti representation of continuous-state branching processes. *Probab. Surveys*, 6:62–89, 2009.

[10] D. A. DAWSON and Z. LI. Stochastic equations, flows and measure-valued processes. *Annals of Probability*, 40(2):813–857, 2012.

[11] T. DUQUESNE and J.-F. LE GALL. Random Trees, Lévy Processes and Spatial Branching Processes. *Astérique*, 281, 2002.

[12] T. DUQUESNE and J.-F. LE GALL. Probabilistic and fractal aspect of Lévy trees. *Probab. Th. Relat. Fields*, 131:553–603, 2005.

[13] Z. F. FU and Z. LI. Stochastic equations of non-negative processes with jumps. *Stochastic Process. Appl.*, 120:306–330, 2010.

[14] K. ITÔ. Poisson point processes attached to Markov processes. *Proceedings Sixth Berkeley Symposium on Mathematical Statistic and Probability*, III:225–239, 1972.

[15] F. B. KNIGHT. Random Walks and A Sojourn Density Process of Brownian Motion . *Transactions of the American Mathematical Society*, 109(1):56–86, 1963.

[16] A. E. KYPRIANOU. *Introductory Lectures on Fluctuations of Lévy Processes with Applications*. Springer-Verlag, Berlin, 2006.

[17] A. LAMBERT. The branching process with logistic growth. *Annals of Applied Probability*, 15(2):1506–1535, 2005.

[18] J. LAMPERTI. Continuous-state branching processes. *Bull. Amer. Math. Soc.*, 73:382–386, 1967.

[19] V. LE, E. PARDOUX, and A. WAKOLBINGER. "Trees under attack": a Ray-Knight representation of Feller's branching diffusion with logistic growth. *Probab. Relat. Fields*, 155(3–4):583–619, 2013.

[20] J.-F. LE GALL and Y. LE JAN. Branching processes in Lévy processes: The exploration process. *Ann. Probab.*, 26:213–252, 1998.

[21] Z. LI. *Measure-valued Branching Markov Processes*. Springer-Verlag, 2011.

[22] E. PARDOUX and A. WAKOLBINGER. From Brownian motion with local time drift to Feller's branching diffusion with logistic growth. *Electric. Comm. Probab.*, 16:720–731, 2011.

[23] E. PARDOUX and A. WAKOLBINGER. From exploration paths to mass excursions - variations on a theme of Ray and Knight. *Surveys in Stochastic Processes, Proceedings of the 33rd SPA Conference in Berlin, 2009*, EMS 2011.

[24] D. RAY. Sojourn times of diffusion processes . *Illinois J. Math.*, 7(4):615–630, 1963.

[25] J.B. WALSH. An introduction to stochastic partial differential equations. In *École d'été de probabilités de Saint-Flour, XIV - 1984*, volume 1180 of *Lecture Notes in Math.*, pages 265–439. Springer, Berlin, 1986.